



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/32, C07K 14/325, A01N 63/00,</b> <b>C12N 15/82</b>	<b>A2</b>	<b>(11) International Publication Number: WO 97/40162</b>  <b>(43) International Publication Date: 30 October 1997 (30.10.97)</b>
<p><b>(21) International Application Number:</b> PCT/US97/06463</p> <p><b>(22) International Filing Date:</b> 18 April 1997 (18.04.97)</p> <p><b>(30) Priority Data:</b> 08/633,993                      19 April 1996 (19.04.96)                      US</p> <p><b>(71) Applicant:</b> MYCOGEN CORPORATION [US/US]; 5501 Oberlin Drive, San Diego, CA 92121 (US).</p> <p><b>(72) Inventors:</b> NARVA, Kenneth, E.; 12123 Caminito Mira Del Mar, San Diego, CA 92130 (US). SCHNEPF, H., Ernest; 7954 Handel Court, San Diego, CA 92126 (US). KNUTH, Mark; 16744 Valle Verde Road, Poway, CA 92074 (US). POLLARD, Michael, R.; 12358 Pathos Lane, San Diego, CA 92129 (US). CARDINEAU, Guy; 17041 Cloudecroft Drive, Poway, CA 92064 (US). SCHWAB, George, E.; 1351 Walnutview, Encinitas, CA 92024 (US).</p> <p><b>(74) Agents:</b> SANDERS, Jay, M. et al.; Saliwanchik, Lloyd &amp; Saliwanchik, P.A., Suite A-1, 2421 N.W. 41st Street, Gainesville, FL 32606-6669 (US).</p>		<p><b>(81) Designated States:</b> AU, CA, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b>  <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p><b>(54) Title:</b> PESTICIDAL TOXINS</p>		
<p><b>(57) Abstract</b></p> <p>The subject invention concerns new classes of pesticidal toxins and polynucleotide sequences which encode these toxins. Also described are novel pesticidal isolates of <i>Bacillus thuringiensis</i>.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## DESCRIPTION

### PESTICIDAL TOXINS

#### Cross-Reference to a Related Application

This application is a continuation-in-part of Application Serial No. 08/633,993, filed April 19, 1996.

#### Background of the Invention

The soil microbe *Bacillus thuringiensis* (*B.t.*) is a Gram-positive, spore-forming bacterium characterized by parasporal crystalline protein inclusions. These inclusions often appear microscopically as distinctively shaped crystals. The proteins can be highly toxic to pests and specific in their toxic activity. Certain *B.t.* toxin genes have been isolated and sequenced, and recombinant DNA-based *B.t.* products have been produced and approved for use. In addition, with the use of genetic engineering techniques, new approaches for delivering these *B.t.* endotoxins to agricultural environments are under development, including the use of plants genetically engineered with endotoxin genes for insect resistance and the use of stabilized intact microbial cells as *B.t.* endotoxin delivery vehicles (Gaertner, F.H., L. Kim [1988] *TIBTECH* 6:S4-S7). Thus, isolated *B.t.* endotoxin genes are becoming commercially valuable.

Until the last ten years, commercial use of *B.t.* pesticides has been largely restricted to a narrow range of lepidopteran (caterpillar) pests. Preparations of the spores and crystals of *B. thuringiensis* subsp. *kurstaki* have been used for many years as commercial insecticides for lepidopteran pests. For example, *B. thuringiensis* var. *kurstaki* HD-1 produces a crystalline  $\delta$ -endotoxin which is toxic to the larvae of a number of lepidopteran insects.

In recent years, however, investigators have discovered *B.t.* pesticides with specificities for a much broader range of pests. For example, other species of *B.t.*, namely *israelensis* and *tenebrionis* (a.k.a. *B.t.* M-7, a.k.a. *B.t. san diego*), have been used commercially to control insects of the orders Diptera and Coleoptera, respectively (Gaertner, F.H. [1989] "Cellular Delivery Systems for Insecticidal Proteins: Living and Non-Living Microorganisms," in *Controlled Delivery of Crop Protection Agents*, R.M. Wilkins, ed., Taylor and Francis, New York and London, 1990, pp. 245-255). See also Couch, T.L. (1980) "Mosquito Pathogenicity of *Bacillus thuringiensis* var. *israelensis*," *Developments in Industrial Microbiology* 22:61-76; Beegle, C.C., (1978) "Use of Entomogenous Bacteria in Agroecosystems," *Developments in Industrial Microbiology* 20:97-104. Krieg, A., A.M. Huger, G.A. Langenbruch, W. Schnetter

(1983) *Z. ang. Ent.* 96:500-508, describe *Bacillus thuringiensis* var. *tenebrionis*, which is reportedly active against two beetles in the order Coleoptera. These are the Colorado potato beetle, *Leptinotarsa decemlineata*, and *Agelastica alni*.

Recently, new subspecies of *B.t.* have been identified, and genes responsible for active  
5  $\delta$ -endotoxin proteins have been isolated (Höfte, H., H.R. Whiteley [1989] *Microbiological Reviews* 52(2):242-255). Höfte and Whiteley classified *B.t.* crystal protein genes into 4 major classes. The classes were CryI (Lepidoptera-specific), CryII (Lepidoptera- and Diptera-specific), CryIII (Coleoptera-specific), and CryIV (Diptera-specific). The discovery of strains specifically toxic to other pests has been reported. (Feitelson, J.S., J. Payne, L. Kim [1992] *Bio/Technology*  
10 10:271-275).

The cloning and expression of a *B.t.* crystal protein gene in *Escherichia coli* has been described in the published literature (Schnepf, H.E., H.R. Whiteley [1981] *Proc. Natl. Acad. Sci. USA* 78:2893-2897). U.S. Patent 4,448,885 and U.S. Patent 4,467,036 both disclose the expression of *B.t.* crystal protein in *E. coli*. U.S. Patents 4,797,276 and 4,853,331 disclose *B.*  
15 *thuringiensis* strain *tenebrionis* (a.k.a. M-7, a.k.a. *B.t. san diego*) which can be used to control coleopteran pests in various environments. U.S. Patent No. 4,918,006 discloses *B.t.* toxins having activity against Dipterans. U.S. Patent No. 4,849,217 discloses *B.t.* isolates which have activity against the alfalfa weevil. U.S. Patent No. 5,208,077 discloses coleopteran-active *Bacillus thuringiensis* isolates. U.S. Patent No. 5,151,363 and U.S. Patent No. 4,948,734  
20 disclose certain isolates of *B.t.* which have activity against nematodes. As a result of extensive research and investment of resources, other patents have issued for new *B.t.* isolates and new uses of *B.t.* isolates. However, the discovery of new *B.t.* isolates and new uses of known *B.t.* isolates remains an empirical, unpredictable art.

Coleopterans are an important group of agricultural pests which cause a very large  
25 amount of damage each year. Examples of coleopteran pests include alfalfa weevils and corn rootworm.

The alfalfa weevil, *Hypera postica*, and the closely related Egyptian alfalfa weevil, *Hypera brunneipennis*, are the most important insect pests of alfalfa grown in the United States, with 2.9 million acres infested in 1984. An annual sum of 20 million dollars is spent to control  
30 these pests. The Egyptian alfalfa weevil is the predominant species in the southwestern U.S., where it undergoes aestivation (*i.e.*, hibernation) during the hot summer months. In all other respects, it is identical to the alfalfa weevil, which predominates throughout the rest of the U.S.

The larval stage is the most damaging in the weevil life cycle. By feeding at the alfalfa plant's growing tips, the larvae cause skeletonization of leaves, stunting, reduced plant growth,

and, ultimately, reductions in yield. Severe infestations can ruin an entire cutting of hay. The adults, also foliar feeders, cause additional, but less significant, damage.

Approximately 9.3 million acres of U.S. corn are infested with corn rootworm species complex each year. The corn rootworm species complex includes the northern corn rootworm, *Diabrotica barberi*, the southern corn rootworm, *D. undecimpunctata howardi*, and the western corn rootworm, *D. virgifera virgifera*. The soil-dwelling larvae of these *Diabrotica* species feed on the root of the corn plant, causing lodging. Lodging eventually reduces corn yield and often results in death of the plant. By feeding on cornsilks, the adult beetles reduce pollination and, therefore, detrimentally effect the yield of corn per plant. In addition, adults and larvae of the genus *Diabrotica* attack cucurbit crops (cucumbers, melons, squash, etc.) and many vegetable and field crops in commercial production as well as those being grown in home gardens.

Control of corn rootworm has been partially addressed by cultivation methods, such as crop rotation and the application of high nitrogen levels to stimulate the growth of an adventitious root system. However, chemical insecticides are relied upon most heavily to guarantee the desired level of control. Insecticides are either banded onto or incorporated into the soil. The major problem associated with the use of chemical insecticides is the development of resistance among the treated insect populations.

#### Brief Summary of the Invention

The subject invention concerns novel materials and methods for controlling non-mammalian pests. In a preferred embodiment, the subject invention provides materials and methods for the control of coleopterian pests. In specific embodiments, the materials and methods described herein are used to control alfalfa weevil and/or corn rootworm.

The subject invention advantageously provides two new classes of polynucleotide sequences which encode corresponding novel classes of pesticidal proteins. One novel class of polynucleotide sequences as described herein encodes toxins which have a full-length molecular weight of approximately 40-50 kDa. In a specific embodiment, these toxins have a molecular weight of about 43-47 kDa. A second class of polynucleotides, which encodes pesticidal proteins of about 10-15 kDa, is also provided according to the subject invention. In a specific embodiment, these toxins have a molecular weight of about 13-14 kDa. The subject invention concerns polynucleotides which encode the 40-50 kDa and 10-15 kDa toxins, polynucleotides which encode pesticidal fragments of the full length toxins, and polynucleotide sequences which encode longer forms of these toxins which include, for example, a protoxin region. In a

preferred embodiment, these toxins, including the fragments, are active against coleopteran pests.

Specific *B.t.* toxins useful according to the invention include toxins which can be obtained from the *B.t.* isolates designated as PS80JJ1, PS149B1, and PS167H2. Of these, PS149B1 and PS167H2 are novel isolates. The subject invention also includes the use of variants of the exemplified *B.t.* isolates and toxins which have substantially the same coleopteran-active properties as the specifically exemplified *B.t.* isolates and toxins. Such variant isolates would include, for example, mutants. Procedures for making mutants are well known in the microbiological art. Ultraviolet light and chemical mutagens such as nitrosoguanidine are used extensively toward this end.

In one embodiment of the subject invention, the polynucleotide sequences of the subject invention are used to encode toxins of approximately 43-47 kDa. These toxins are then used to control coleopteran pests. In a particularly preferred embodiment, the coleopteran pests are corn rootworms. The genes which encode the 43-47 kDa toxins can be obtained from, for example, PS80JJ1, PS149B1, or PS167H2. In a second embodiment, toxins of approximately 13-14 kDa are used to control coleopteran pests. The approximately 13-14 kDa toxin, as well as the genes which encode these toxins, can also be obtained from PS80JJ1, PS149B1, or PS167H2. In a particularly preferred embodiment, the activity of the 43-47 kDa toxins can be augmented and/or facilitated by further contacting the target pests with an approximately 13-14 kDa toxin.

In a preferred embodiment, the subject invention concerns plants cells transformed with at least one polynucleotide sequence of the subject invention such that the transformed plant cells express pesticidal toxins in tissues consumed by the target pests.

Alternatively, the *B.t.* isolates of the subject invention, or recombinant microbes expressing the toxins described herein, can be used to control pests. In this regard, the invention includes the treatment of substantially intact *B.t.* cells, and/or recombinant cells containing the expressed toxins of the invention, treated to prolong the pesticidal activity when the substantially intact cells are applied to the environment of a target pest. The treated cell acts as a protective coating for the pesticidal toxin. The toxin becomes active upon ingestion by a target insect.

#### Brief Description of the Drawings

**Figure 1** shows three specific 43-47 kDa pesticidal toxins of the subject invention as well as a consensus sequence for these pesticidal toxins.

**Figure 2** shows the relationship of the 14 and 45 kDa sequences of PS80JJ1 (SEQ ID NOS. 31 and 10).

Brief Description of the Sequences

SEQ ID NO. 1 is a 5-amino acid N-terminal sequence of the approximately 45 kDa toxin of 80JJ1.

SEQ ID NO. 2 is a 25-amino acid N-terminal sequence of the approximately 45 kDa toxin of 80JJ1.

SEQ ID NO. 3 is a 24-amino acid N-terminal sequence of the approximately 14 kDa toxin of 80JJ1.

SEQ ID NO. 4 is the N-terminal sequence of the approximately 47 kDa toxin from 149B1.

SEQ ID NO. 5 is a 50-amino acid N-terminal amino acid sequence for the purified approximately 14 kDa protein from PS149B1.

SEQ ID NO. 6 is the N-terminal sequence of the approximately 47 kDa toxin from 167H2.

SEQ ID NO. 7 is a 25-amino acid N-terminal sequence for the purified approximately 14 kDa protein from PS167H2.

SEQ ID NO. 8 is an oligonucleotide probe for the gene encoding the PS80JJ1 44.3 kDa toxin and is a forward primer for PS149B1 and PS167H2 used according to the subject invention.

SEQ ID NO. 9 is a reverse primer for PS149B1 and PS167H2 used according to the subject invention.

SEQ ID NO. 10 is the nucleotide sequence of the gene encoding the approximately 45 kDa PS80JJ1 toxin.

SEQ ID NO. 11 is the amino acid sequence for the approximately 45 kDa PS80JJ1 toxin.

SEQ ID NO. 12 is the partial nucleotide sequence of the gene encoding the approximately 44 kDa PS149B1 toxin.

SEQ ID NO. 13 is the partial amino acid sequence for the approximately 44 kDa PS149B1 toxin.

SEQ ID NO. 14 is the partial nucleotide sequence of the gene encoding the approximately 44 kDa PS167H2 toxin.

SEQ ID NO. 15 is the partial amino acid sequence for the approximately 44 kDa PS167H2 toxin.

SEQ ID NO. 16 is a peptide sequence used in primer design according to the subject invention.

SEQ ID NO. 17 is a peptide sequence used in primer design according to the subject invention.

SEQ ID NO. 18 is a peptide sequence used in primer design according to the subject invention.

5        SEQ ID NO. 19 is a peptide sequence used in primer design according to the subject invention.

SEQ ID NO. 20 is a nucleotide sequence corresponding to the peptide of SEQ ID NO. 16.

10        SEQ ID NO. 21 is a nucleotide sequence corresponding to the peptide of SEQ ID NO. 17.

SEQ ID NO. 22 is a nucleotide sequence corresponding to the peptide of SEQ ID NO. 18.

SEQ ID NO. 23 is a nucleotide sequence corresponding to the peptide of SEQ ID NO. 19.

15        SEQ ID NO. 24 is a reverse primer based on the reverse complement of SEQ ID NO. 22.

SEQ ID NO. 25 is a reverse primer based on the reverse complement of SEQ ID NO. 23.

SEQ ID NO. 26 is a forward primer based on the PS80JJ1 44.3 kDa toxin.

20        SEQ ID NO. 27 is a reverse primer based on the PS80JJ1 44.3 kDa toxin.

SEQ ID NO. 28 is a generic sequence representing a new class of toxins according to the subject invention.

SEQ ID NO. 29 is an oligonucleotide probe used according to the subject invention.

25        SEQ ID NO. 30 is the nucleotide sequence of the entire genetic locus containing open reading frames of both the 14 and 45 kDa PS80JJ1 toxins and the flanking nucleotide sequences.

SEQ ID NO. 31 is the nucleotide sequence of the PS80JJ1 14 kDa toxin open reading frame.

SEQ ID NO. 32 is the deduced amino acid sequence of the 14 kDa toxin of PS80JJ1.

30        SEQ ID NO. 33 is a reverse oligonucleotide primer used according to the subject invention.

SEQ ID NO. 34 is the nucleotide sequence of the entire genetic locus containing open reading frames of both the 14 and 44 kDa PS167H2 toxins and the flanking nucleotide sequences.

SEQ ID NO. 35 is the nucleotide sequence of the gene encoding the approximately 14 kDa PS167H2 toxin.

SEQ ID NO. 36 is the amino acid sequence for the approximately 14 kDa PS167H2 toxin.

5 SEQ ID NO. 37 is the nucleotide sequence of the gene encoding the approximately 44 kDa PS167H2 toxin.

SEQ ID NO. 38 is the amino acid sequence for the approximately 44 kDa PS167H2 toxin.

10 SEQ ID NO. 39 is the nucleotide sequence of the entire genetic locus containing open reading frames of both the 14 and 44 kDa PS149B1 toxins and the flanking nucleotide sequences.

SEQ ID NO. 40 is the nucleotide sequence of the gene encoding the approximately 14 kDa PS149B1 toxin.

15 SEQ ID NO. 41 is the amino acid sequence for the approximately 14 kDa PS149B1 toxin.

SEQ ID NO. 42 is the nucleotide sequence of the gene encoding the approximately 44 kDa PS149B1 toxin.

SEQ ID NO. 43 is the amino acid sequence for the approximately 44 kDa PS149B1 toxin.

20 SEQ ID NO. 44 is a maize-optimized gene sequence encoding the approximately 14 kDa toxin of 80JJ1.

SEQ ID NO. 45 is a maize-optimized gene sequence encoding the approximately 44 kDa toxin of 80JJ1.

25 Detailed Disclosure of the Invention

The subject invention concerns two new classes of polynucleotide sequences which encode novel pesticidal toxins. In one embodiment, the toxins have a full-length molecular weight of approximately 40-50 kDa. In specific embodiments exemplified herein, these toxins have a molecular weight of about 43-47 kDa. In a second embodiment, the pesticidal toxins have a molecular weight of approximately 10-15 kDa. In specific embodiments exemplified herein, these toxins have a molecular weight of about 13-14 kDa. Certain specific toxins are exemplified herein. For toxins having a known amino acid sequence, the molecular weight is also known. Those skilled in the art will recognize that the apparent molecular weight of a protein as determined by gel electrophoresis will sometimes differ from the true molecular

30

weight. Therefore, reference herein to, for example, a 45 kDa protein or a 14 kDa protein is understood to refer to proteins of approximately that size even if the true molecular weight is somewhat different.

The subject invention concerns not only the polynucleotide sequences which encode these classes of toxins, but also the use of these polynucleotide sequences to produce recombinant hosts which express the toxins. In a further aspect, the subject invention concerns the combined use of an approximately 40-50 kDa toxin of the subject invention together with an approximately 10-15 kDa toxin of the subject invention to achieve highly effective control of pests, including coleopterans such as corn rootworm.

A further aspect of the subject invention concerns two novel isolates and the toxins and genes obtainable from these isolates. The novel *B.t.* isolates of the subject invention have been designated PS149B1 and PS167H2.

The new classes of toxins and polynucleotide sequences provided here are defined according to several parameters. One critical characteristic of the toxins described herein is pesticidal activity. In a specific embodiment, these toxins have activity against coleopteran pests. The toxins and genes of the subject invention can be further defined by their amino acid and nucleotide sequences. The sequences of the molecules within each novel class can be defined herein in terms of homology to certain exemplified sequences as well as in terms of the ability to hybridize with, or be amplified by, certain exemplified probes and primers. The classes of toxins provided herein can also be identified based on their immunoreactivity with certain antibodies and based upon their adherence to a generic formula.

The sequence of three approximately 45 kDa toxins of the subject invention are provided as SEQ ID NOS. 11, 43, and 38. In a preferred embodiment of the subject invention, the toxins in this new class have a sequence which conforms to the generic sequence presented as SEQ ID NO. 28. In a specific embodiment, the toxins of this class will conform to the consensus sequence shown in Figure 1.

In a preferred embodiment, the toxins of the subject invention have at least one of the following characteristics:

- (a) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO. 4, DNA which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which encodes SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO. 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes

SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25, SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38, SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43;

- (b) said toxin immunoreacts with an antibody to an approximately 40-50 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554;
- (c) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using a primer pair selected from the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp;
- (d) said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28;
- (e) said toxin comprises an amino acid sequence which has at least about 60% homology with a pesticidal portion of an amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43;
- (f) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO. 5, DNA which encodes SEQ ID NO. 7, DNA which encodes SEQ ID NO. 32, DNA which encodes SEQ ID NO. 36, and DNA which encodes SEQ ID NO. 41;
- (g) said toxin immunoreacts with an antibody to an approximately 10-15 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying

characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554;

- (h) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using the primer pair of SEQ ID NO. 29 and SEQ ID NO. 33; and
- (i) said toxin comprises an amino acid sequence which has at least about 60% homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of SEQ ID NO. 41.

As used herein "stringent" conditions for hybridization refers to conditions which achieve the same, or about the same, degree of specificity of hybridization as the conditions employed by the current applicants. Specifically, hybridization of immobilized DNA on Southern blots with <sup>32</sup>P-labeled gene-specific probes was performed by standard methods (Maniatis, T., E.F. Fritsch, J. Sambrook [1982] *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.). In general, hybridization and subsequent washes were carried out under stringent conditions that allowed for detection of target sequences with homology to the PS80JJ1 toxin genes. For double-stranded DNA gene probes, hybridization was carried out overnight at 20-25° C below the melting temperature (T<sub>m</sub>) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula (Beltz, G.A., K.A. Jacobs, T.H. Eickbush, P.T. Cherbas, and F.C. Kafatos [1983] *Methods of Enzymology*, R. Wu, L. Grossman and K. Moldave [eds.] Academic Press, New York 100:266-285).

$$T_m = 81.5^{\circ} \text{C} + 16.6 \log[\text{Na}^+] + 0.41(\% \text{G+C}) - 0.61(\% \text{formamide}) - 600/\text{length of duplex in base pairs}.$$

Washes are typically carried out as follows:

- (1) Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at T<sub>m</sub>-20°C for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

For oligonucleotide probes, hybridization was carried out overnight at 10-20°C below the melting temperature (T<sub>m</sub>) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. T<sub>m</sub> for oligonucleotide probes was determined by the following formula:

$$T_m (^{\circ}C) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$$

(Suggs, S.V., T. Miyake, E.H. Kawashime, M.J. Johnson, K. Itakura, and R.B. Wallace [1981] *ICN-UCLA Symp. Dev. Biol. Using Purified Genes*, D.D. Brown [ed.], Academic Press, New York, 23:683-693).

5 Washes were typically carried out as follows:

- (1) Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

10 With the teachings provided herein, one skilled in the art could readily produce and use the various toxins and polynucleotide sequences of the novel classes described herein.

Microorganisms useful according to the subject invention have been deposited in the permanent collection of the Agricultural Research Service Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 North University Street, Peoria, Illinois 61604, USA.

15 The culture repository numbers of the deposited strains are as follows:

<u>Culture</u>	<u>Repository No.</u>	<u>Deposit Date</u>
<i>B.t.</i> strain PS80JJ1	NRRL B-18679	July 17, 1990
<i>B.t.</i> strain PS149B1	NRRL B-21553	March 28, 1996
<i>B.t.</i> strain PS167H2	NRRL B-21554	March 28, 1996
20 <i>E. coli</i> NM522 (pMYC2365)	NRRL B-21170	January 5, 1994
<i>E. coli</i> NM522 (pMYC2382)	NRRL B-21329	September 28, 1994
<i>E. coli</i> NM522 (pMYC2379)	NRRL B-21155	November 3, 1993
<i>E. coli</i> NM522(pMYC2421)	NRRL B-21555	March 28, 1996
<i>E. coli</i> NM522(pMYC2427)	NRRL B-21672	March 26, 1997
25 <i>E. coli</i> NM522(pMYC2429)	NRRL B-21673	March 26, 1997
<i>E. coli</i> NM522(pMYC2426)	NRRL B-21671	March 26, 1997

The PS80JJ1 isolate is available to the public by virtue of the issuance of U.S. Patent No. 5,151,363.

30 *B.t.* isolates PS149B1 and PS167H2 have been deposited under conditions that assure that access to the cultures will be available during the pendency of this patent application to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 CFR 1.14 and 35 U.S.C. 122. The deposits are available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny, are filed. However, it

should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

Further, the subject culture deposits will be stored and made available to the public in accord with the provisions of the Budapest Treaty for the Deposit of Microorganisms, i.e., they will be stored with all the care necessary to keep them viable and uncontaminated for a period of at least five years after the most recent request for the furnishing of a sample of a deposit, and in any case, for a period of at least 30 (thirty) years after the date of deposit or for the enforceable life of any patent which may issue disclosing the cultures. The depositor acknowledges the duty to replace the deposit(s) should the depository be unable to furnish a sample when requested, due to the condition of the deposit(s). All restrictions on the availability to the public of the subject culture deposits will be irrevocably removed upon the granting of a patent disclosing them.

Following is a table which provides characteristics of certain *B.t.* isolates useful according to the subject invention.

**Table 1.** Description of *B.t.* strains toxic to coleopterans

Culture	Crystal Description	Approx. MW (kDa)	Serotype	NRRL Deposit	Deposit Date
PS80JJ1	multiple attached	130, 90, 47, 37, 14	4a4b, sotto	B-18679	7-17-90
PS149B1		130, 47, 14		B-21553	3-28-96
PS167H2		70, 47, 14		B-23554	3-28-96

**Genes and toxins.** The genes and toxins useful according to the subject invention include not only the full length sequences disclosed but also fragments of these sequences, variants, mutants, and fusion proteins which retain the characteristic pesticidal activity of the toxins specifically exemplified herein. As used herein, the terms "variants" or "variations" of genes refer to nucleotide sequences which encode the same toxins or which encode equivalent toxins having pesticidal activity. As used herein, the term "equivalent toxins" refers to toxins having the same or essentially the same biological activity against the target pests as the claimed toxins.

It should be apparent to a person skilled in this art that genes encoding active toxins can be identified and obtained through several means. The specific genes exemplified herein may be obtained from the isolates deposited at a culture depository as described above. These genes, or portions or variants thereof, may also be constructed synthetically, for example, by use of a gene synthesizer. Variations of genes may be readily constructed using standard techniques for making point mutations. Also, fragments of these genes can be made using commercially available exonucleases or endonucleases according to standard procedures. For example, enzymes such as *Bal31* or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes. Also, genes which encode active fragments may be obtained using a variety of restriction enzymes. Proteases may be used to directly obtain active fragments of these toxins.

Equivalent toxins and/or genes encoding these equivalent toxins can be derived from *B.t.* isolates and/or DNA libraries using the teachings provided herein. There are a number of methods for obtaining the pesticidal toxins of the instant invention. For example, antibodies to the pesticidal toxins disclosed and claimed herein can be used to identify and isolate other toxins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the toxins which are most constant and most distinct from other *B.t.* toxins. These antibodies can then be used to specifically identify equivalent toxins with the characteristic activity by immunoprecipitation, enzyme linked immunosorbent assay (ELISA), or western blotting. Antibodies to the toxins disclosed herein, or to equivalent toxins, or fragments of these toxins, can readily be prepared using standard procedures in this art. The genes which encode these toxins can then be obtained from the microorganism.

Fragments and equivalents which retain the pesticidal activity of the exemplified toxins would be within the scope of the subject invention. Also, because of the redundancy of the genetic code, a variety of different DNA sequences can encode the amino acid sequences disclosed herein. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding the same, or essentially the same, toxins. These variant DNA sequences are within the scope of the subject invention. As used herein, reference to "essentially the same" sequence refers to sequences which have amino acid substitutions, deletions, additions, or insertions which do not materially affect pesticidal activity. Fragments retaining pesticidal activity are also included in this definition.

A further method for identifying the toxins and genes of the subject invention is through the use of oligonucleotide probes. These probes are detectable nucleotide sequences. These sequences may be detectable by virtue of an appropriate label or may be made inherently

fluorescent as described in International Application No. WO93/16094. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong bond between the two molecules, it can be reasonably assumed that the probe and sample have substantial homology. Preferably, hybridization is conducted under stringent conditions by techniques well-known in the art, as described, for example, in Keller, G.H., M.M. Manak (1987) *DNA Probes*, Stockton Press, New York, NY., pp. 169-170. Detection of the probe provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying toxin-encoding genes of the subject invention. The nucleotide segments which are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures. These nucleotide sequences can also be used as PCR primers to amplify genes of the subject invention.

Certain toxins of the subject invention have been specifically exemplified herein. Since these toxins are merely exemplary of the toxins of the subject invention, it should be readily apparent that the subject invention comprises variant or equivalent toxins (and nucleotide sequences coding for equivalent toxins) having the same or similar pesticidal activity of the exemplified toxin. Equivalent toxins will have amino acid homology with an exemplified toxin. The amino acid identity will typically be greater than 60%, preferably be greater than 75%, more preferably greater than 80%, more preferably greater than 90%, and can be greater than 95%. The amino acid homology will be highest in critical regions of the toxin which account for biological activity or are involved in the determination of three-dimensional configuration which ultimately is responsible for the biological activity. In this regard, certain amino acid substitutions are acceptable and can be expected if these substitutions are in regions which are not critical to activity or are conservative amino acid substitutions which do not affect the three-dimensional configuration of the molecule. For example, amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the subject invention so long as the substitution does not materially alter the biological activity of the compound. Table 2 provides a listing of examples of amino acids belonging to each class.

Table 2.

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not significantly detract from the biological activity of the toxin.

The toxins of the subject invention can also be characterized in terms of the shape and location of toxin inclusions, which are described above.

Recombinant hosts. The toxin-encoding genes harbored by the isolates of the subject invention can be introduced into a wide variety of microbial or plant hosts. Expression of the toxin gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide. With suitable microbial hosts, *e.g.*, *Pseudomonas*, the microbes can be applied to the situs of the pest, where they will proliferate and be ingested. The result is a control of the pest. Alternatively, the microbe hosting the toxin gene can be treated under conditions that prolong the activity of the toxin and stabilize the cell. The treated cell, which retains the toxic activity, then can be applied to the environment of the target pest.

Where the *B.t.* toxin gene is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, it is essential that certain host microbes be used. Microorganism hosts are selected which are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplane) of one or more crops of interest. These microorganisms are selected so as to be capable of successfully competing in the particular environment (crop and other insect habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

A large number of microorganisms are known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, *e.g.*, genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*,

*Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, e.g., genera *Saccharomyces*, *Cryptococcus*, *Cluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacterium tumefaciens*, *Rhodopseudomonas spheroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes entrophus*, and *Azotobacter vinlandii*; and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odoratus*, *Cluyveromyces veronae*, and *Aureobasidium pullulans*. Of particular interest are the pigmented microorganisms.

A wide variety of ways are available for introducing a *B.t.* gene encoding a toxin into a microorganism host under conditions which allow for stable maintenance and expression of the gene. These methods are well known to those skilled in the art and are described, for example, in United States Patent No. 5,135,867, which is incorporated herein by reference.

Control of coleopterans, including corn rootworm using the isolates, toxins, and genes of the subject invention can be accomplished by a variety of methods known to those skilled in the art. These methods include, for example, the application of *B.t.* isolates to the pests (or their location), the application of recombinant microbes to the pests (or their locations), and the transformation of plants with genes which encode the pesticidal toxins of the subject invention. Recombinant microbes may be, for example, a *B.t.*, *E. coli*, or *Pseudomonas*. Transformations can be made by those skilled in the art using standard techniques. Materials necessary for these transformations are disclosed herein or are otherwise readily available to the skilled artisan.

Synthetic genes which are functionally equivalent to the toxins of the subject invention can also be used to transform hosts. Methods for the production of synthetic genes can be found in, for example, U.S. Patent No. 5,380,831.

Control of other pests such as lepidopterans and other insects, nematodes, and mites can also be accomplished by those skilled in the art using standard techniques combined with the teachings provided herein.

Treatment of cells. As mentioned above, *B.t.* or recombinant cells expressing a *B.t.* toxin can be treated to prolong the toxin activity and stabilize the cell. The pesticide microcapsule that is formed comprises the *B.t.* toxin within a cellular structure that has been stabilized and will protect the toxin when the microcapsule is applied to the environment of the target pest. Suitable host cells may include either prokaryotes or eukaryotes, normally being

limited to those cells which do not produce substances toxic to higher organisms, such as mammals. However, organisms which produce substances toxic to higher organisms could be used, where the toxic substances are unstable or the level of application sufficiently low as to avoid any possibility of toxicity to a mammalian host. As hosts, of particular interest will be the prokaryotes and the lower eukaryotes, such as fungi.

The cell will usually be intact and be substantially in the proliferative form when treated, rather than in a spore form, although in some instances spores may be employed.

Treatment of the microbial cell, e.g., a microbe containing the *B.t.* toxin gene, can be by chemical or physical means, or by a combination of chemical and/or physical means, so long as the technique does not deleteriously affect the properties of the toxin, nor diminish the cellular capability of protecting the toxin. Examples of chemical reagents are halogenating agents, particularly halogens of atomic no. 17-80. More particularly, iodine can be used under mild conditions and for sufficient time to achieve the desired results. Other suitable techniques include treatment with aldehydes, such as glutaraldehyde; anti-infectives, such as zephiran chloride and cetylpyridinium chloride; alcohols, such as isopropyl and ethanol; various histologic fixatives, such as Lugol iodine, Bouin's fixative, various acids and Helly's fixative (See: Humason, Gretchen L., *Animal Tissue Techniques*, W.H. Freeman and Company, 1967); or a combination of physical (heat) and chemical agents that preserve and prolong the activity of the toxin produced in the cell when the cell is administered to the host environment. Examples of physical means are short wavelength radiation such as gamma-radiation and X-radiation, freezing, UV irradiation, lyophilization, and the like. Methods for treatment of microbial cells are disclosed in United States Patent Nos. 4,695,455 and 4,695,462, which are incorporated herein by reference.

The cells generally will have enhanced structural stability which will enhance resistance to environmental conditions. Where the pesticide is in a proform, the method of cell treatment should be selected so as not to inhibit processing of the proform to the mature form of the pesticide by the target pest pathogen. For example, formaldehyde will crosslink proteins and could inhibit processing of the proform of a polypeptide pesticide. The method of treatment should retain at least a substantial portion of the bio-availability or bioactivity of the toxin.

Characteristics of particular interest in selecting a host cell for purposes of production include ease of introducing the *B.t.* gene into the host, availability of expression systems, efficiency of expression, stability of the pesticide in the host, and the presence of auxiliary genetic capabilities. Characteristics of interest for use as a pesticide microcapsule include protective qualities for the pesticide, such as thick cell walls, pigmentation, and intracellular

packaging or formation of inclusion bodies; survival in aqueous environments; lack of mammalian toxicity; attractiveness to pests for ingestion; ease of killing and fixing without damage to the toxin; and the like. Other considerations include ease of formulation and handling, economics, storage stability, and the like.

5           Growth of cells. The cellular host containing the *B.t.* insecticidal gene may be grown in any convenient nutrient medium, where the DNA construct provides a selective advantage, providing for a selective medium so that substantially all or all of the cells retain the *B.t.* gene. These cells may then be harvested in accordance with conventional ways. Alternatively, the cells can be treated prior to harvesting.

10           The *B.t.* cells of the invention can be cultured using standard art media and fermentation techniques. Upon completion of the fermentation cycle the bacteria can be harvested by first separating the *B.t.* spores and crystals from the fermentation broth by means well known in the art. The recovered *B.t.* spores and crystals can be formulated into a wettable powder, liquid concentrate, granules or other formulations by the addition of surfactants, dispersants, inert carriers, and other components to facilitate handling and application for particular target pests. 15           These formulations and application procedures are all well known in the art.

Formulations. Formulated bait granules containing an attractant and spores and crystals of the *B.t.* isolates, or recombinant microbes comprising the genes obtainable from the *B.t.* isolates disclosed herein, can be applied to the soil. Formulated product can also be applied as 20           a seed-coating or root treatment or total plant treatment at later stages of the crop cycle. Plant and soil treatments of *B.t.* cells may be employed as wettable powders, granules or dusts, by mixing with various inert materials, such as inorganic minerals (phyllsilicates, carbonates, sulfates, phosphates, and the like) or botanical materials (powdered corncobs, rice hulls, walnut shells, and the like). The formulations may include spreader-sticker adjuvants, stabilizing 25           agents, other pesticidal additives, or surfactants. Liquid formulations may be aqueous-based or non-aqueous and employed as foams, gels, suspensions, emulsifiable concentrates, or the like. The ingredients may include rheological agents, surfactants, emulsifiers, dispersants, or polymers.

30           As would be appreciated by a person skilled in the art, the pesticidal concentration will vary widely depending upon the nature of the particular formulation, particularly whether it is a concentrate or to be used directly. The pesticide will be present in at least 1% by weight and may be 100% by weight. The dry formulations will have from about 1-95% by weight of the pesticide while the liquid formulations will generally be from about 1-60% by weight of the solids in the liquid phase. The formulations will generally have from about  $10^2$  to about  $10^4$

cells/mg. These formulations will be administered at about 50 mg (liquid or dry) to 1 kg or more per hectare.

The formulations can be applied to the environment of the pest, *e.g.*, soil and foliage, by spraying, dusting, sprinkling, or the like.

5           Mutants. Mutants of the isolates of the invention can be made by procedures well known in the art. For example, an asporogenous mutant can be obtained through ethylmethane sulfonate (EMS) mutagenesis of an isolate. The mutants can be made using ultraviolet light and nitrosoguanidine by procedures well known in the art.

10           A smaller percentage of the asporogenous mutants will remain intact and not lyse for extended fermentation periods; these strains are designated lysis minus (-). Lysis minus strains can be identified by screening asporogenous mutants in shake flask media and selecting those mutants that are still intact and contain toxin crystals at the end of the fermentation. Lysis minus strains are suitable for a cell treatment process that will yield a protected, encapsulated toxin protein.

15           To prepare a phage resistant variant of said asporogenous mutant, an aliquot of the phage lysate is spread onto nutrient agar and allowed to dry. An aliquot of the phage sensitive bacterial strain is then plated directly over the dried lysate and allowed to dry. The plates are incubated at 30°C. The plates are incubated for 2 days and, at that time, numerous colonies could be seen growing on the agar. Some of these colonies are picked and subcultured onto  
20           nutrient agar plates. These apparent resistant cultures are tested for resistance by cross streaking with the phage lysate. A line of the phage lysate is streaked on the plate and allowed to dry. The presumptive resistant cultures are then streaked across the phage line. Resistant bacterial cultures show no lysis anywhere in the streak across the phage line after overnight incubation at 30°C. The resistance to phage is then reconfirmed by plating a lawn of the resistant culture  
25           onto a nutrient agar plate. The sensitive strain is also plated in the same manner to serve as the positive control. After drying, a drop of the phage lysate is placed in the center of the plate and allowed to dry. Resistant cultures showed no lysis in the area where the phage lysate has been placed after incubation at 30°C for 24 hours.

30           Following are examples which illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

Example 1 – Culturing of *B.t.* Isolates of the Invention

A subculture of the *B.t.* isolates, or mutants thereof, can be used to inoculate the following medium, a peptone, glucose, salts medium.

5	Bacto Peptone	7.5 g/l
	Glucose	1.0 g/l
	$\text{KH}_2\text{PO}_4$	3.4 g/l
	$\text{K}_2\text{HPO}_4$	4.35 g/l
	Salt Solution	5.0 ml/l
10	$\text{CaCl}_2$ Solution	5.0 ml/l
	pH 7.2	
	Salts Solution (100 ml)	
	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	2.46 g
15	$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	0.04 g
	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.28 g
	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.40 g
	$\text{CaCl}_2$ Solution (100 ml)	
20	$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	3.66 g

The salts solution and  $\text{CaCl}_2$  solution are filter-sterilized and added to the autoclaved and cooked broth at the time of inoculation. Flasks are incubated at 30°C on a rotary shaker at 200 rpm for 64 hr.

25 The above procedure can be readily scaled up to large fermentors by procedures well known in the art.

The *B.t.* spores and/or crystals, obtained in the above fermentation, can be isolated by procedures well known in the art. A frequently-used procedure is to subject the harvested fermentation broth to separation techniques, e.g., centrifugation.

30

Example 2 – Protein Purification for 45 kDa 80JJ1 Protein

One gram of lyophilized powder of 80JJ1 was suspended in 40 ml of buffer containing 80 mM Tris-Cl pH 7.8, 5 mM EDTA, 100  $\mu\text{M}$  PMSF, 0.5  $\mu\text{g/ml}$  Leupeptin, 0.7  $\mu\text{g/ml}$  Pepstatin, and 40  $\mu\text{g/ml}$  Bestatin. The suspension was centrifuged, and the resulting supernatant was

discarded. The pellet was washed five times using 35-40 ml of the above buffer for each wash. The washed pellet was resuspended in 10 ml of 40% NaBr, 5 mM EDTA, 100  $\mu$ M PMSF, 0.5  $\mu$ g/ml Leupeptin, 0.7  $\mu$ g/ml Pepstatin, and 40  $\mu$ g/ml Bestatin and placed on a rocker platform for 75 minutes. The NaBr suspension was centrifuged, the supernatant was removed, and the pellet was treated a second time with 40% NaBr, 5 mM EDTA, 100  $\mu$ M PMSF, 0.5  $\mu$ g/ml Leupeptin, 0.7  $\mu$ g/ml Pepstatin, and 40  $\mu$ g/ml Bestatin as above. The supernatants (40% NaBr soluble) were combined and dialyzed against 10 mM CAPS pH 10.0, 1 mM EDTA at 4°C. The dialyzed extracts were centrifuged and the resulting supernatant was removed. The pellet (40% NaBr dialysis pellet) was suspended in 5 ml of H<sub>2</sub>O and centrifuged. The resultant supernatant was removed and discarded. The washed pellet was washed a second time in 10 ml of H<sub>2</sub>O and centrifuged as above. The washed pellet was suspended in 1.5 ml of H<sub>2</sub>O and contained primarily three peptides with molecular weights of approximately 47 kDa, 45 kDa, and 15 kDa when analyzed using SDS-PAGE. At this stage of purification, the suspended 40% NaBr dialysis pellet contained approximately 21 mg/ml of protein by Lowry assay.

The peptides in the pellet suspension were separated using SDS-PAGE (Laemmli, U.K. [1970] *Nature* 227:680) in 15% acrylamide gels. The separated proteins were then electrophoretically blotted to a PVDF membrane (Millipore Corp.) in 10 mM CAPS pH 11.0, 10% MeOH at 100 V constant. After one hour the PVDF membrane was rinsed in water briefly and placed for 3 minutes in 0.25% Coomassie blue R-250, 50% methanol, 5% acetic acid. The stained membrane was destained in 40% MeOH, 5% acetic acid. The destained membrane was air-dried at room temperature (LeGendre *et al.* [1989] In *A Practical Guide to Protein Purification For Microsequencing*, P. Matsudaira, ed., Academic Press, New York, NY). The membrane was sequenced using automated gas phase Edman degradation (Hunkapillar, M.W., R.M. Hewick, W.L. Dreyer, L.E. Hood [1983] *Meth. Enzymol.* 91:399).

The amino acid analysis revealed that the N-terminal sequence of the 45 kDa band was as follows: Met-Leu-Asp-Thr-Asn (SEQ ID NO. 1).

The 47 kDa band was also analyzed and the N-terminal amino acid sequence was determined to be the same 5-amino acid sequence as SEQ ID NO. 1. Therefore, the N-terminal amino acid sequences of the 47 kDa peptide and the 45 kDa peptide were identical.

This amino acid sequence also corresponds to a sequence obtained from a 45 kDa peptide obtained from PS80JJ1 spore/crystal powders, using another purification protocol, with the N-terminal sequence as follows: Met-Leu-Asp-Thr-Asn-Lys-Val-Tyr-Glu-Ile-Ser-Asn-Leu-Ala-Asn-Gly-Leu-Tyr-Thr-Ser-Thr-Tyr-Leu-Ser-Leu (SEQ ID NO. 2).

Example 3 – Purification of the 14 kDa Peptide of PS80III

0.8 ml of the white dialysis suspension (approximately 21 mg/ml) containing the 47 kDa, 45 kDa, and 15 kDa peptides, was dissolved in 10 ml of 40% NaBr, and 0.5 ml of 100 mM EDTA were added. After about 18 hours (overnight), a white opaque suspension was obtained. This was collected by centrifugation and discarded. The supernatant was concentrated in a Centricon-30 (Amicon Corporation) to a final volume of approximately 15 ml. The filtered volume was washed with water by filter dialysis and incubated on ice, eventually forming a milky white suspension. The suspension was centrifuged and the pellet and supernatant were separated and retained. The pellet was then suspended in 1.0 ml water (approximately 6 mg/ml). The pellet contained substantially pure 15 kDa protein when analyzed by SDS-PAGE.

The N-terminal amino acid sequence was determined to be: Ser-Ala-Arg-Glu-Val-His-Ile-Glu-Ile-Asn-Asn-Thr-Arg-His-Thr-Leu-Gln-Leu-Glu-Ala-Lys-Thr-Lys-Leu (SEQ ID NO. 3).

Example 4 – Protein Purification and Characterization of PS149B1 45 kDa Protein

The P1 pellet was resuspended with two volumes of deionized water per unit wet weight, and to this was added nine volumes of 40% (w/w) aqueous sodium bromide. This and all subsequent operations were carried out on ice or at 4-6°C. After 30 minutes, the suspension was diluted with 36 volumes of chilled water and centrifuged at 25,000 x g for 30 minutes to give a pellet and a supernatant.

The resulting pellet was resuspended in 1-2 volumes of water and layered on a 20-40% (w/w) sodium bromide gradient and centrifuged at 8,000 x g for 100 minutes. The layer banding at approximately 32% (w/w) sodium bromide (the "inclusions", or INC) was recovered and dialyzed overnight against water using a dialysis membrane with a 6-8 kDa MW cut-off. Particulate material was recovered by centrifugation at 25,000 x g, resuspended in water, and aliquoted and assayed for protein by the method of Lowry and by SDS-PAGE.

The resulting supernatant was concentrated 3- to 4-fold using Centricon-10 concentrators, then dialyzed overnight against water using a dialysis membrane with a 6-8 kDa MW cut-off. Particulate material was recovered by centrifugation at 25,000 x g, resuspended in water, and aliquoted and assayed for protein by the method of Lowry and by SDS-PAGE. This fraction was denoted as P1.P2.

The peptides in the pellet suspension were separated using SDS-PAGE (Laemlli, U.K., *supra*) in 15% acrylamide gels. The separated proteins were then electrophoretically blotted to a PVDF membrane (Millipore Corp.) in 10 mM CAPS pH 11.0, 10% MeOH at 100 V constant.

After one hour the PVDF membrane was rinsed in water briefly and placed for 3 minutes in 0.25% Coomassie blue R-250, 50% methanol, 5% acetic acid. The stained membrane was destained in 40% MeOH, 5% acetic acid. The destained membrane was air-dried at room temperature (LeGendre *et al.*, *supra*). The membrane was sequenced using automated gas phase Edman degradation (Hunkapillar *et al.*, *supra*).

Protein analysis indicated the presence of two major polypeptides, with molecular weights of 47 kDa and 14 kDa. Molecular weights were measured against standard polypeptides of known molecular weight. This process provides only an estimate of true molecular weight. The 47 kDa band from PS149B1 migrated on SDS-PAGE in a manner indistinguishable from the 47 kDa protein from PS80JJ1. Likewise, the 14 kDa band from PS149B1 migrated on SDS-PAGE in a manner indistinguishable from 14 kDa bands from PS167H2 and PS80JJ1. Apart from these two polypeptides, which were estimated to account for 25-35% (47 kDa) and 35-55% (15 kDa) of the Coomassie staining material respectively, there may be minor bands, including those of estimated MW at 46 kDa, 130 kDa, and 70 kDa.

Protein analysis indicated that fraction INC contained a single polypeptide with MW of 47 kDa, and that fraction P1.P2 contained a single polypeptide with MW of 14 kDa. These polypeptides were recovered in yields greater than 50% from P1.

The N-terminal amino acid sequence for the purified 47 kDa protein from PS149B1 is: Met-Leu-Asp-Thr-Asn-Lys-Val-Tyr-Glu-Ile-Ser-Asn-His-Ala-Asn-Gly-Leu-Tyr-Ala-Ala-Thr-Tyr-Leu-Ser-Leu (SEQ ID NO. 4).

The N-terminal amino acid sequence for the purified 14 kDa protein from PS149B1 is: Ser-Ala-Arg-Glu-Val-His-Ile-Asp-Val-Asn-Asn-Lys-Thr-Gly-His-Thr-Leu-Gln-Leu-Glu-Asp-Lys-Thr-Lys-Leu-Asp-Gly-Gly-Arg-Trp-Arg-Thr-Ser-Pro-Xaa-Asn-Val-Ala-Asn-Asp-Gln-Ile-Lys-Thr-Phe-Val-Ala-Glu-Ser-Asn (SEQ ID NO. 5).

#### Example 5 – Amino Acid Sequence for 45 kDa and 14 kDa Toxins of PS167H2

The N-terminal amino acid sequence for the purified 45 kDa protein from PS167H2 is: Met-Leu-Asp-Thr-Asn-Lys-Ile-Tyr-Glu-Ile-Ser-Asn-Tyr-Ala-Asn-Gly-Leu-His-Ala-Ala-Thr-Tyr-Leu-Ser-Leu (SEQ ID NO. 6).

The N-terminal amino acid sequence for the purified 14 kDa protein from PS167H2 is: Ser-Ala-Arg-Glu-Val-His-Ile-Asp-Val-Asn-Asn-Lys-Thr-Gly-His-Thr-Leu-Gln-Leu-Glu-Asp-Lys-Thr-Lys-Leu (SEQ ID NO. 7).

These amino acid sequences can be compared to the sequence obtained for the 47 kDa peptide obtained from 80JJ1 spore/crystal powders with the N-terminal sequence (SEQ ID NO.

1) and to the sequence obtained for the 14 kDa peptide obtained from 80JJ1 spore/crystal powders with the N-terminal sequence (SEQ ID NO. 3).

Clearly, the 45-47 kDa proteins are highly related and probably represent one gene family, and the 14 kDa proteins are highly related and probably represent another gene family.

5        Example 6 – Molecular Cloning, Expression, and DNA Sequence Analysis of a Novel  $\delta$ -Endotoxin Gene from *Bacillus thuringiensis* Strain PS80JJ1

10        Total cellular DNA was prepared from *Bacillus thuringiensis* (B.t.) cells grown to an optical density, at 600 nm, of 1.0. Cells were pelleted by centrifugation and resuspended in protoplast buffer (20 mg/ml lysozyme in 0.3 M sucrose, 25 mM Tris-Cl [pH 8.0], 25 mM EDTA). After incubation at 37°C for 1 hour, protoplasts were lysed by two cycles of freezing and thawing. Nine volumes of a solution of 0.1 M NaCl, 0.1% SDS, 0.1 M Tris-Cl were added to complete lysis. The cleared lysate was extracted twice with phenol:chloroform (1:1). Nucleic acids were precipitated with two volumes of ethanol and pelleted by centrifugation. The pellet  
15        was resuspended in TE buffer and RNase was added to a final concentration of 50 µg/ml. After incubation at 37°C for 1 hour, the solution was extracted once each with phenol:chloroform (1:1) and TE-saturated chloroform. DNA was precipitated from the aqueous phase by the addition of one-tenth volume of 3 M NaOAc and two volumes of ethanol. DNA was pelleted by centrifugation, washed with 70% ethanol, dried, and resuspended in TE buffer.

20        An oligonucleotide probe for the gene encoding the PS80JJ1 45 kDa toxin was designed from N-terminal peptide sequence data. The sequence of the 29-base oligonucleotide probe was: 5'-ATG YTW GAT ACW AAT AAA GTW TAT GAA AT-3' (SEQ ID NO. 8)

This oligonucleotide was mixed at four positions as shown. This probe was radiolabeled with <sup>32</sup>P and used in standard condition hybridization of Southern blots of PS80JJ1 total cellular DNA  
25        digested with various restriction endonucleases. Representative autoradiographic data from these experiments showing the sizes of DNA restriction fragments containing sequence homology to the 44.3 kDa toxin oligonucleotide probe of SEQ ID NO. 8 are presented in Table 3.

**Table 3.** RFLP of PS80JJ1 cellular DNA fragments on Southern blots that hybridized under standard conditions with the 44.3 kDa toxin gene oligonucleotide probe (SEQ ID NO. 8)

Restriction Enzyme	Approximate Fragment Size (kbp)
<i>EcoRI</i>	6.0
<i>HindIII</i>	8.3
<i>KpnI</i>	7.4
<i>PstI</i>	11.5
<i>XbaI</i>	9.1

These DNA fragments identified in these analyses contain all or a segment of the PS80JJ1 45 kDa toxin gene. The approximate sizes of the hybridizing DNA fragments in Table 3 are in reasonable agreement with the sizes of a subset of the PS80JJ1 fragments hybridizing with a PS80JJ1 45 kDa toxin subgene probe used in separate experiments, as predicted (see Table 4, below).

A gene library was constructed from PS80JJ1 DNA partially digested with *Sau3AI*. Partial restriction digests were fractionated by agarose gel electrophoresis. DNA fragments 9.3 to 23 kbp in size were excised from the gel, electroeluted from the gel slice, purified on an Elutip-D ion exchange column (Schleicher and Schuell, Keene, NH), and recovered by ethanol precipitation. The *Sau3AI* inserts were ligated into *Bam*HI-digested LambdaGem-11 (Promega, Madison, WI). Recombinant phage were packaged and plated on *E. coli* KW251 cells. Plaques were screened by hybridization with the oligonucleotide probe described above. Hybridizing phage were plaque-purified and used to infect liquid cultures of *E. coli* KW251 cells for isolation of DNA by standard procedures (Maniatis *et al.*, *supra*).

Southern blot analysis revealed that one of the recombinant phage isolates contained an approximately 4.8 kbp *XbaI*-*SacI* band that hybridized to the PS80JJ1 toxin gene probe. The *SacI* site flanking the PS80JJ1 toxin gene is a phage vector cloning site, while the flanking *XbaI* site is located within the PS80JJ1 DNA insert. This DNA restriction fragment was subcloned by standard methods into pBluescript S/K (Stratagene, San Diego, CA) for sequence analysis. The resultant plasmid was designated pMYC2421. The DNA insert was also subcloned into pHTBlueII (an *E. coli/B. thuringiensis* shuttle vector comprised of pBluescript S/K [Stratagene, La Jolla, CA] and the replication origin from a resident *B.t.* plasmid [D. Lereclus *et al.* (1989) *FEMS Microbiology Letters* 60:211-218]) to yield pMYC2420.

An oligonucleotide probe for the gene encoding the PS80JJ1 14 kDa toxin was designed from N-terminal peptide sequence data. The sequence of the 28-base oligonucleotide probe was: 5' GW GAA GTW CAT ATW GAA ATW AAT AAT AC 3' (SEQ ID NO. 29). This oligonucleotide was mixed at four positions as shown. The probe was radiolabelled with <sup>32</sup>P and used in standard condition hybridizations of Southern blots of PS80JJ1 total cellular and pMYC2421 DNA digested with various restriction endonucleases. These RFLP mapping experiments demonstrated that the gene encoding the 14 kDa toxin is located on the same genomic *EcoRI* fragment that contains the N-terminal coding sequence for the 44.3 kDa toxin.

To test expression of the PS80JJ1 toxin genes in *B.t.*, pMYC2420 was transformed into the acrySTALLIFEROUS (Cry-) *B.t.* host, CryB (A. Aronson, Purdue University, West Lafayette, IN), by electroporation. Expression of both the approximately 14 and 44.3 kDa PS80JJ1 toxins encoded by pMYC2420 was demonstrated by SDS-PAGE analysis. Toxin crystal preparations from the recombinant CryB[pMYC2420] strain, MR536, were assayed and found to be active against western corn rootworm.

The PS80JJ1 toxin genes encoded by pMYC2421 were sequenced using the ABI373 automated sequencing system and associated software. The sequence of the entire genetic locus containing both open reading frames and flanking nucleotide sequences is shown in SEQ ID NO. 30. The termination codon of the 14 kDa toxin gene is 121 base pairs upstream (5') from the initiation codon of the 44.3 kDa toxin gene (Figure 2). The PS80JJ1 14 kDa toxin open reading frame nucleotide sequence (SEQ ID NO. 31), the 44.3 kDa toxin open reading frame nucleotide sequence (SEQ ID NO. 10), and the respective deduced amino acid sequences (SEQ ID NO. 32 and SEQ ID NO. 11) are novel compared to other toxin genes encoding pesticidal proteins.

Thus, the nucleotide sequence encoding the 14 kDa toxin of PS80JJ1 is shown in SEQ ID NO. 31. The deduced amino acid sequence of the 14 kDa toxin of PS80JJ1 is shown in SEQ ID NO. 32. The nucleotide sequences encoding both the 14 and 45 kDa toxins of PS80JJ1, as well as the flanking sequences, are shown in SEQ ID NO. 30. The relationship of these sequences is shown in Figure 2.

A subculture of *E. coli* NM522 containing plasmid pMYC2421 was deposited in the permanent collection of the Patent Culture Collection (NRRL), Regional Research Center, 1815 North University Street, Peoria, IL 61604 USA on March 28, 1996. The accession number is NRRL B-21555.

Example 7 – RFLP and PCR Analysis of Additional Novel  $\delta$ -Endotoxin Genes from *Bacillus thuringiensis* Strains PS149B1 and PS167H2

Two additional strains active against corn rootworm, PS149B1 and PS167H2, also produce parasporal protein crystals comprised in part of polypeptides approximately 14 and 45 kDa in size. Southern hybridization and partial DNA sequence analysis were used to examine the relatedness of these toxins to the 80JJ1 toxins. DNA was extracted from these *B.t.* strains as described above, and standard Southern hybridizations were performed using the 14 kDa toxin oligonucleotide probe (SEQ ID NO. 29) and an approximately 800 bp PCR fragment of the 80JJ1 44.3 kDa toxin gene-encoding sequence. Representative RFLP data from these experiments showing the sizes of DNA restriction fragments containing sequence homology to the 44.3 kDa toxin are presented in Table 4. Representative RFLP data from these experiments showing the sizes of DNA restriction fragments containing sequence homology to the approximately 14 kDa toxin are presented in Table 5.

**Table 4.** RFLP of PS80JJ1, PS149B1, and PS167H2 cellular DNA fragments on Southern blots that hybridized with the approximately 800 bp PS80JJ1 44.3 kDa toxin subgene probe under standard conditions

Restriction enzyme	Strain		
	PS80JJ1	PS149B1	PS167H2
Approximate fragment size (kbp)			
<i>EcoRI</i>	6.4	5.7	2.6
	1.3	2.8	
	0.6		
<i>HindIII</i>	8.2	6.2	4.4
<i>KpnI</i>	7.8	10.0	11.5
<i>PstI</i>	12.0	9.2	9.2
			8.2
<i>XbaI</i>	9.4	10.9	10.9
<i>SacI</i>	17.5	15.5	11.1
	13.1	10.5	6.3

Each of the three strains exhibited unique RFLP patterns. The hybridizing DNA fragments from PS149B1 or PS167H2 contain all or part of toxin genes with sequence homology to the PS80JJ1 44.3 kDa toxin.

**Table 5.** Restriction fragment length polymorphisms of PS80JJ1, PS149B1, and PS167H2 cellular DNA fragments on Southern blots that hybridized with the PS80JJ1 14 kDa toxin oligonucleotide probe under standard conditions

Restriction enzyme	Strain		
	PS80JJ1	PS149B1	PS167H2
Approximate fragment size (kbp)			
<i>EcoRI</i>	5.6	2.7	2.7
<i>HindIII</i>	7.1	6.0	4.7
<i>XbaI</i>	8.4	11.2	11.2

Each of the three strains exhibited unique RFLP patterns. The hybridizing DNA fragments from PS149B1 or PS167H2 contain all or part of toxin genes with sequence homology to the PS80JJ1 14 kDa toxin gene.

Portions of the toxin genes in PS149B1 or PS167H2 were amplified by PCR using forward and reverse oligonucleotide primer pairs designed based on the PS80JJ1 44.3 kDa toxin gene sequence. For PS149B1, the following primer pair was used:

Forward:

5'-ATG YTW GAT ACW AAT AAA GTW TAT GAA AT-3' (SEQ ID NO. 8)

Reverse:

5'-GGA TTA TCT ATC TCT GAG TGT TCT TG-3' (SEQ ID NO. 9)

For PS167H2, the same primer pair was used. These PCR-derived fragments were sequenced using the ABI373 automated sequencing system and associated software. The partial gene and peptide sequences obtained are shown in SEQ ID NO. 12-15. These sequences contain portions of the nucleotide coding sequences and peptide sequences for novel corn rootworm-active toxins present in *B.t.* strains PS149B1 or PS167H2.

Example 8 – Molecular Cloning and DNA Sequence Analysis of Novel  $\delta$ -Endotoxin Genes from *Bacillus thuringiensis* Strains PS149B1 and PS167H2

Total cellular DNA was extracted from strains PS149B1 and PS167H2 as described for PS80JJ1. Gene libraries of size-fractionated *Sau*3A partial restriction fragments were constructed in Lambda-Gem11 for each respective strain as previously described. Recombinant phage were packaged and plated on *E. coli* KW251 cells. Plaques were screened by hybridization with the oligonucleotide probe specific for the 44 kDa toxin gene. Hybridizing phage were plaque-purified and used to infect liquid cultures of *E. coli* KW251 cells for isolation of DNA by standard procedures (Maniatis *et al.*, *supra*).

For PS167H2, Southern blot analysis revealed that one of the recombinant phage isolates contained an approximately 4.0 to 4.4 kbp *Hind*III band that hybridized to the PS80JJ1 44 kDa toxin gene 5' oligonucleotide probe (SEQ ID NO. 8). This DNA restriction fragment was subcloned by standard methods into pBluescript S/K (Stratgene, San Diego, CA) for sequence analysis. The fragment was also subcloned into the high copy number shuttle vector, pHT370 (Arantes, O., D. Lereclus [1991] *Gene* 108:115-119) for expression analyses in *Bacillus thuringiensis* (see below). The resultant recombinant, high copy number bifunctional plasmid was designated pMYC2427.

The PS167H2 toxin genes encoded by pMYC2427 were sequenced using the ABI automated sequencing system and associated software. The sequence of the entire genetic locus containing both open reading frames and flanking nucleotide sequences is shown in SEQ ID NO. 34. The termination codon of the 14 kDa toxin gene is 107 base pairs upstream (5') from the initiation codon of the 44 kDa toxin gene. The PS167H2 14 kDa toxin coding sequence (SEQ ID NO. 35), the 44 kDa toxin coding sequence (SEQ ID NO. 37), and the respective deduced amino acid sequences, SEQ ID NO. 36 and SEQ ID NO. 38, are novel compared to other known toxin genes encoding pesticidal proteins. The toxin genes are arranged in a similar manner to, and have some homology with, the PS80JJ1 14 and 44 kDa toxins.

A subculture of *E. coli* NM522 containing plasmid pMYC2427 was deposited in the permanent collection of the Patent Culture Collection (NRRL), Regional Research Center, 1815 North University Street, Peoria, Illinois 61604 USA on 26 March 1997. The accession number is NRRL B-21672.

For PS149B1, Southern blot analysis using the PS80JJ1 44 kDa oligonucleotide 5' probe (SEQ ID NO. 8) demonstrated hybridization of an approximately 5.9 kbp *Cla*I DNA fragment. Complete *Cla*I digests of PS149B1 genomic DNA were size fractionated on agarose gels and cloned into pHTBlueII. The fragment was also subcloned into the high copy number shuttle

vector, pHT370 (Arantes, O., D. Lereclus [1991] *Gene* 108:115-119) for expression analyses in *Bacillus thuringiensis* (see below). The resultant recombinant, high copy number bifunctional plasmid was designated pMYC2429.

The PS149B1 toxin genes encoded by pMYC2429 were sequenced using the ABI automated sequencing system and associated software. The sequence of the entire genetic locus containing both open reading frames and flanking nucleotide sequences is shown in SEQ ID NO. 39. The termination codon of the 14 kDa toxin gene is 108 base pairs upstream (5') from the initiation codon of the 44 kDa toxin gene. The PS149B1 14 kDa toxin coding sequence (SEQ ID NO. 40), the 44 kDa toxin coding sequence (SEQ ID NO. 42), and the respective deduced amino acid sequences, SEQ ID NO. 41 and SEQ ID NO. 43, are novel compared to other known toxin genes encoding pesticidal proteins. The toxin genes are arranged in a similar manner as, and have some homology with, the PS80JJ1 and PS167H2 14 and 44 kDa toxins. Together, these three toxin operons comprise a new family of pesticidal toxins.

A subculture of *E. coli* NM522 containing plasmid pMYC2429 was deposited in the permanent collection of the Patent Culture Collection (NRRL), Regional Research Center, 1815 North University Street, Peoria, Illinois 61604 USA on 26 March 1997. The accession number is NRRL B-21673.

#### Example 9 – PCR Amplification for Identification and Cloning Novel Corn Rootworm-Active Toxin

The DNA and peptide sequences of the three novel approximately 45 kDa corn rootworm-active toxins from PS80JJ1, PS149B1, and PS167H2 (SEQ ID NOS. 12-15) were aligned with the Genetics Computer Group sequence analysis program Pileup using a gap weight of 3.00 and a gap length weight of 0.10. The sequence alignments were used to identify conserved peptide sequences to which oligonucleotide primers were designed that were likely to hybridize to genes encoding members of this novel toxin family. Such primers can be used in PCR to amplify diagnostic DNA fragments for these and related toxin genes. Numerous primer designs to various sequences are possible, four of which are described here to provide an example. These peptide sequences are:

Asp-Ile-Asp-Asp-Tyr-Asn-Leu (SEQ ID NO. 16)

Trp-Phe-Leu-Phe-Pro-Ile-Asp (SEQ ID NO. 17)

Gln-Ile-Lys-Thr-Thr-Pro-Tyr-Tyr (SEQ ID NO. 18)

Tyr-Glu-Trp-Gly-Thr-Glu (SEQ ID NO. 19).

The corresponding nucleotide sequences are:

5'-GATATWGATGAYTAYAAYTTR-3' (SEQ ID NO. 20)

5'-TGGTTTTTTRTTTCCWATWGAY-3' (SEQ ID NO. 21)

5'-CAAATHAAACWACWCCATATTAT-3' (SEQ ID NO. 22)

5'-TAYGARTGGGGHACAGAA-3' (SEQ ID NO. 23).

Forward primers for polymerase amplification in thermocycle reactions were designed based on the nucleotide sequences of SEQ ID NOS. 20 and 21.

Reverse primers were designed based on the reverse complement of SEQ ID NOS. 22 and 23:

5'-ATAATATGGWGTGTTTDDATTG-3' (SEQ ID NO. 24)

5'-TTCTGTDCCCCACTCRTA-3' (SEQ ID NO. 25).

These primers can be used in combination to amplify DNA fragments of the following sizes (Table 6) that identify genes encoding novel corn rootworm toxins.

**Table 6.** Predicted sizes of diagnostic DNA fragments (base pairs) amplifiable with primers specific for novel corn rootworm-active toxins

Primer pair (SEQ ID NO.)	DNA fragment size (bp)
20 + 24	495
20 + 25	594
21 + 24	471
21 + 25	580

Similarly, entire genes encoding novel corn rootworm-active toxins can be isolated by polymerase amplification in thermocycle reactions using primers designed based on DNA sequences flanking the open reading frames. For the PS80JJ1 44.3 kDa toxin, one such primer pair was designed, synthesized, and used to amplify a diagnostic 1613 bp DNA fragment that included the entire toxin coding sequence. These primers are:

Forward: 5'-CTCAAAGCGGATCAGGAG-3' (SEQ ID NO. 26)

Reverse: 5'-GCGTATTCGGATATGCTTGG-3' (SEQ ID NO. 27).

For PCR amplification of the PS80JJ1 14 kDa toxin, the oligonucleotide coding for the N-terminal peptide sequence (SEQ ID NO. 29) can be used in combination with various reverse oligonucleotide primers based on the sequences in the PS80JJ1 toxin gene locus. One such reverse primer has the following sequence:

5' CATGAGATTATCTCCTGATCCGC 3' (SEQ ID NO. 33).

When used in standard PCR reactions, this primer pair amplified a diagnostic 1390 bp DNA fragment that includes the entire 14 kDa toxin coding sequence and some 3' flanking sequences corresponding to the 121 base intergenic spacer and a portion of the 44.3 kDa toxin gene. When used in combination with the 14 kDa forward primer, PCR will generate a diagnostic 322 base pair DNA fragment.

#### Example 10 – Bioassay of Protein

A preparation of the insoluble fraction from the dialyzed NaBr extract of 80JJ1 containing the 47 kDa, 45 kDa, and 15 kDa peptides was bioassayed against Western corn rootworm and found to exhibit significant toxin activity.

#### Example 11 – Bioassay of Protein

The purified protein fractions from PS149B1 were bioassayed against western corn rootworm and found to exhibit significant toxin activity when combined. In fact, the combination restored activity to that noted in the original preparation (P1). The following bioassay data set presents percent mortality and demonstrates this effect.

**Table 7.**

Concentration ( $\mu\text{g}/\text{cm}^2$ )	P1	INC	P1.P2	INC + P1.P2
300	88, 100, 94	19	13	100
100	94, 50, 63	31	38	94
33.3	19, 19, 44	38	13	50
11.1	13, 56, 25	12	31	13
3.7	0, 50, 0	0	31	13
1.2	13, 43, 12	0	12	19
0.4	6, 12, 6	25	19	6

#### Example 12 – Clone Dose-Response Bioassays

The PS80JJ1 toxin operon was also subcloned from pMYC2421 into pHT370 for direct comparison of bioactivity with the recombinant toxins cloned from PS149B1 and PS167H2. The resultant recombinant, high copy number bifunctional plasmid was designated pMYC2426.

A subculture of *E. coli* NM522 containing plasmid pMYC2426 was deposited in the permanent collection of the Patent Culture Collection (NRRL), Regional Research Center, 1815 North University Street, Peoria, Illinois 61604 USA on 26 March 1997. The accession number is NRRL B-21671.

To test expression of the PS80JJ1, PS149B1 and PS167H2 toxin genes in *B.t.*, pMYC2426, pMYC2427 and pMYC2429 were separately transformed into the acrySTALLIFEROUS (Cry-) *B. t.* host, CryB (A. Aronson, Purdue University, West Lafayette, IN), by electroporation. The recombinant strains were designated MR543 (CryB [pMYC2426]), MR544 (CryB [pMYC2427]) and MR546 (CryB [pMYC2429]), respectively. Expression of both the approximately 14 and 44 kDa toxins was demonstrated by SDS-PAGE analysis for each recombinant strain.

Toxin crystal preparations from the recombinant strains were assayed against western corn rootworm. Their diet was amended with sorbic acid and SIGMA pen-strep-ampho-B. The material was top-loaded at a rate of 50  $\mu$ l of suspension per  $\text{cm}^2$  diet surface area. Bioassays were run with neonate Western corn rootworm larvae for 4 days at approximately 25°C. Percentage mortality and top-load  $\text{LC}_{50}$  estimates for the clones (pellets) are set forth in Table 8.

**Table 8.**

Sample	Percentage mortality at given protein concentration ( $\mu\text{g}/\text{cm}^2$ )		
	50	5	0.5
MR543 pellet	44	19	9
MR544 pellet	72	32	21
MR546 pellet	52	32	21
dH <sub>2</sub> O	7		

#### Example 13 – Insertion and Expression of Toxin Genes Into Plants

One aspect of the subject invention is the transformation of plants with genes encoding the insecticidal toxin. The transformed plants are resistant to attack by the target pest.

The novel corn rootworm-active genes described here can be optimized for expression in other organisms. Maize optimized gene sequences encoding the 14 and 44 kDa PS80JJ1 toxins are disclosed in SEQ ID NO. 44 and SEQ ID NO. 45, respectively.

Genes encoding pesticidal toxins, as disclosed herein, can be inserted into plant cells using a variety of techniques which are well known in the art. For example, a large number of cloning vectors comprising a replication system in *E. coli* and a marker that permits selection of the transformed cells are available for preparation for the insertion of foreign genes into higher plants. The vectors comprise, for example, pBR322, pUC series, M13mp series, pACYC184, etc. Accordingly, the sequence encoding the *B.t.* toxin can be inserted into the vector at a suitable restriction site. The resulting plasmid is used for transformation into *E. coli*. The *E. coli* cells are cultivated in a suitable nutrient medium, then harvested and lysed. The plasmid is recovered. Sequence analysis, restriction analysis, electrophoresis, and other biochemical-molecular biological methods are generally carried out as methods of analysis. After each manipulation, the DNA sequence used can be cleaved and joined to the next DNA sequence. Each plasmid sequence can be cloned in the same or other plasmids. Depending on the method of inserting desired genes into the plant, other DNA sequences may be necessary. If, for example, the Ti or Ri plasmid is used for the transformation of the plant cell, then at least the right border, but often the right and the left border of the Ti or Ri plasmid T-DNA, has to be joined as the flanking region of the genes to be inserted.

The use of T-DNA for the transformation of plant cells has been intensively researched and sufficiently described in EP 120 516; Hoekema (1985) In: *The Binary Plant Vector System*, Offset-durkkerij Kanters B.V., Alblaserdam, Chapter 5; Fraley *et al.*, *Crit. Rev. Plant Sci.* 4:1-46; and An *et al.* (1985) *EMBO J.* 4:277-287.

Once the inserted DNA has been integrated in the genome, it is relatively stable there and, as a rule, does not come out again. It normally contains a selection marker that confers on the transformed plant cells resistance to a biocide or an antibiotic, such as kanamycin, G 418, bleomycin, hygromycin, or chloramphenicol, *inter alia*. The individually employed marker should accordingly permit the selection of transformed cells rather than cells that do not contain the inserted DNA.

A large number of techniques are available for inserting DNA into a plant host cell. Those techniques include transformation with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation agent, fusion, injection, biolistics (microparticle bombardment), or electroporation as well as other possible methods. If *Agrobacteria* are used for the transformation, the DNA to be inserted has to be cloned into special plasmids, namely either into an intermediate vector or into a binary vector. The intermediate vectors can be integrated into the Ti or Ri plasmid by homologous recombination owing to sequences that are homologous to sequences in the T-DNA. The Ti or Ri plasmid also comprises the vir region

necessary for the transfer of the T-DNA. Intermediate vectors cannot replicate themselves in *Agrobacteria*. The intermediate vector can be transferred into *Agrobacterium tumefaciens* by means of a helper plasmid (conjugation). Binary vectors can replicate themselves both in *E. coli* and in *Agrobacteria*. They comprise a selection marker gene and a linker or polylinker which are framed by the right and left T-DNA border regions. They can be transformed directly into *Agrobacteria* (Holsters *et al.* [1978] *Mol. Gen. Genet.* 163:181-187). The *Agrobacterium* used as host cell is to comprise a plasmid carrying a vir region. The vir region is necessary for the transfer of the T-DNA into the plant cell. Additional T-DNA may be contained. The bacterium so transformed is used for the transformation of plant cells. Plant explants can advantageously be cultivated with *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* for the transfer of the DNA into the plant cell. Whole plants can then be regenerated from the infected plant material (for example, pieces of leaf, segments of stalk, roots, but also protoplasts or suspension-cultivated cells) in a suitable medium, which may contain antibiotics or biocides for selection. The plants so obtained can then be tested for the presence of the inserted DNA. No special demands are made of the plasmids in the case of injection and electroporation. It is possible to use ordinary plasmids, such as, for example, pUC derivatives.

The transformed cells grow inside the plants in the usual manner. They can form germ cells and transmit the transformed trait(s) to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

In a preferred embodiment of the subject invention, plants will be transformed with genes wherein the codon usage has been optimized for plants. See, for example, U.S. Patent No. 5,380,831, which is hereby incorporated by reference. Also, advantageously, plants encoding a truncated toxin will be used. The truncated toxin typically will encode about 55% to about 80% of the full length toxin. Methods for creating synthetic *B.t.* genes for use in plants are known in the art.

#### Example 14 – Cloning of *B.t.* Genes Into Insect Viruses

A number of viruses are known to infect insects. These viruses include, for example, baculoviruses and entomopoxviruses. In one embodiment of the subject invention, genes encoding the insecticidal toxins, as described herein, can be placed within the genome of the insect virus, thus enhancing the pathogenicity of the virus. Methods for constructing insect viruses which comprise *B.t.* toxin genes are well known and readily practiced by those skilled

in the art. These procedures are described, for example, in Merryweather *et al.* (Merryweather, A.T., U. Weyer, M.P.G. Harris, M. Hirst, T. Booth, R.D. Possee (1990) *J. Gen. Virol.* 71:1535-1544) and Martens *et al.* (Martens, J.W.M., G. Honee, D. Zuidema, J.W.M. van Lent, B. Visser, J.M. Vlak (1990) *Appl. Environmental Microbiol.* 56(9):2764-2770).

5

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and the scope of the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

Applicant Name(s): MYCOGEN CORPORATION  
Street address: 5501 Oberlin Drive  
City: San Diego  
State/Province: California  
Country: US  
Postal code/Zip: 92121  
Phone number: (619) 453-8030 Fax number: (619) 453-6991

## (ii) TITLE OF INVENTION: Pesticidal Toxins

## (iii) NUMBER OF SEQUENCES: 45

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Saliwanchik, Lloyd & Saliwanchik  
(B) STREET: 2421 N.W. 41st Street, Suite A-1  
(C) CITY: Gainesville  
(D) STATE: FL  
(E) COUNTRY: USA  
(F) ZIP: 32606-6669

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US  
(B) FILING DATE:  
(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/633,993  
(B) FILING DATE: 19-APR-1996  
(C) CLASSIFICATION:

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Sanders, Jay M.  
(B) REGISTRATION NUMBER: 39,355  
(C) REFERENCE/DOCKET NUMBER: MA-703C1

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 352-375-8100  
(B) TELEFAX: 352-372-5800

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids  
(B) TYPE: amino acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Leu Asp Thr Asn

1 5

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Leu Asp Thr Asn Lys Val Tyr Glu Ile Ser Asn Leu Ala Asn Gly

1 5 10 15

Leu Tyr Thr Ser Thr Tyr Leu Ser Leu

20 25

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Ser Ala Arg Glu Val His Ile Glu Ile Asn Asn Thr Arg His Thr Leu

1 5 10 15

Gln Leu Glu Ala Lys Thr Lys Leu

20

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```
Met Leu Asp Thr Asn Lys Val Tyr Glu Ile Ser Asn His Ala Asn Gly
 1             5             10             15

Leu Tyr Ala Ala Thr Tyr Leu Ser Leu
          20             25
```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```
Ser Ala Arg Glu Val His Ile Asp Val Asn Asn Lys Thr Gly His Thr
 1             5             10             15

Leu Gln Leu Glu Asp Lys Thr Lys Leu Asp Gly Gly Arg Trp Arg Thr
          20             25             30

Ser Pro Xaa Asn Val Ala Asn Asp Gln Ile Lys Thr Phe Val Ala Glu
          35             40             45

Ser Asn
          50
```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```
Met Leu Asp Thr Asn Lys Ile Tyr Glu Ile Ser Asn Tyr Ala Asn Gly
 1             5             10             15

Leu His Ala Ala Thr Tyr Leu Ser Leu
          20             25
```

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ser	Ala	Arg	Glu	Val	His	Ile	Asp	Val	Asn	Asn	Lys	Thr	Gly	His	Thr
1				5					10					15	
Leu	Gln	Leu	Glu	Asp	Lys	Thr	Lys	Leu							
			20					25							

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATGNTNGATA CNAATAAAGT NTATGAAAT

29

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGATTATCTA TCTCTGAGTG TTCTTG

26

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1158 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATGTTAGATA CTAATAAAGT TTATGAAATA AGCAATCTTG CTAATGGATT ATATACATCA	60
ACTTATTTAA GTCTTGATGA TTCAGGTGTT AGTTTAATGA GTAAAAAGGA TGAAGATATT	120
GATGATTACA ATTTAAATG GTTTTATTT CCTATTGATA ATAATCAATA TATTATTACA	180
AGCTATGGAG CTAATAATTG TAAAGTTTG AATGTTAAAA ATGATAAAAT AAATGTTTCA	240
ACTTATTCTT CAACAACTC TGTACAAAA TGGCAAATAA AAGCTAAAGA TTCTTCATAT	300
ATAATACAAA GTGATAATGG AAAGTCTTA ACAGCAGGAG TAGGTCAATC TCTTGGATA	360
GTACGCCTAA CTGATGAATT TCCAGAGAAT TCTAACCAAC AATGGAATTT AACTCCTGTA	420
CAAACAATTC AACTCCCACA AAAACCTAAA ATAGATGAAA AATTAAAGA TCATCCTGAA	480
TATTCAGAAA CCGGAAATAT AAATCCTAAA ACAACTCCTC AATTAATGGG ATGGACATTA	540
GTACCTTGTA TTATGGTAAA TGATTCAAAA ATAGATAAAA ACACCTCAAT TAAAACCTACT	600
CCATATTATA TTTTAAAAA ATATAAATAC TGGAACTAG CAAAAGGAAG TAATGTATCT	660
TTACTTCCAC ATCAAAAAAG ATCATATGAT TATGAATGGG GTACAGAAAA AAATCAAAAA	720
ACAACTATTA TTAATACAGT AGGATTGCAA ATTAATATAG ATTCAGGAAT GAAATTTGAA	780
GTACCAGAAG TAGGAGGAGG TACAGAAGAC ATAAAAACAC AATTAACCTGA AGAATTAATA	840
GTTGAATATA GCACTGAAAC CAAAATAATG ACGAAATATC AAGAACACTC AGAGATAGAT	900
AATCCAACCT ATCAACCAAT GAATTCCTATA GGACTTCTTA TTTATACTTC TTTAGAATTA	960
TATCGATATA ACGGTACAGA AATTAAGATA ATGGACATAG AAACCTCAGA TCATGATACT	1020
TACACTCTTA CTCTTATCC AAATCATAAA GAAGCATTAT TACTTCTCAC AAACCATTCG	1080
TATGAAGAAG TAGAAGAAAT AACAAAAATA CCTAAGCATA CACTTATAAA ATTGAAAAAA	1140
CATTATTTTA AAAAATAA	1158

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 385 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met	Leu	Asp	Thr	Asn	Lys	Val	Tyr	Glu	Ile	Ser	Asn	Leu	Ala	Asn	Gly
1				5				10					15		

Leu Tyr Thr Ser Thr Tyr Leu Ser Leu Asp Asp Ser Gly Val Ser Leu  
 20 25 30  
 Met Ser Lys Lys Asp Glu Asp Ile Asp Asp Tyr Asn Leu Lys Trp Phe  
 35 40 45  
 Leu Phe Pro Ile Asp Asn Asn Gln Tyr Ile Ile Thr Ser Tyr Gly Ala  
 50 55 60  
 Asn Asn Cys Lys Val Trp Asn Val Lys Asn Asp Lys Ile Asn Val Ser  
 65 70 75 80  
 Thr Tyr Ser Ser Thr Asn Ser Val Gln Lys Trp Gln Ile Lys Ala Lys  
 85 90 95  
 Asp Ser Ser Tyr Ile Ile Gln Ser Asp Asn Gly Lys Val Leu Thr Ala  
 100 105 110  
 Gly Val Gly Gln Ser Leu Gly Ile Val Arg Leu Thr Asp Glu Phe Pro  
 115 120 125  
 Glu Asn Ser Asn Gln Gln Trp Asn Leu Thr Pro Val Gln Thr Ile Gln  
 130 135 140  
 Leu Pro Gln Lys Pro Lys Ile Asp Glu Lys Leu Lys Asp His Pro Glu  
 145 150 155 160  
 Tyr Ser Glu Thr Gly Asn Ile Asn Pro Lys Thr Thr Pro Gln Leu Met  
 165 170 175  
 Gly Trp Thr Leu Val Pro Cys Ile Met Val Asn Asp Ser Lys Ile Asp  
 180 185 190  
 Lys Asn Thr Gln Ile Lys Thr Thr Pro Tyr Tyr Ile Phe Lys Lys Tyr  
 195 200 205  
 Lys Tyr Trp Asn Leu Ala Lys Gly Ser Asn Val Ser Leu Leu Pro His  
 210 215 220  
 Gln Lys Arg Ser Tyr Asp Tyr Glu Trp Gly Thr Glu Lys Asn Gln Lys  
 225 230 235 240  
 Thr Thr Ile Ile Asn Thr Val Gly Leu Gln Ile Asn Ile Asp Ser Gly  
 245 250 255  
 Met Lys Phe Glu Val Pro Glu Val Gly Gly Gly Thr Glu Asp Ile Lys  
 260 265 270  
 Thr Gln Leu Thr Glu Glu Leu Lys Val Glu Tyr Ser Thr Glu Thr Lys  
 275 280 285  
 Ile Met Thr Lys Tyr Gln Glu His Ser Glu Ile Asp Asn Pro Thr Asn  
 290 295 300  
 Gln Pro Met Asn Ser Ile Gly Leu Leu Ile Tyr Thr Ser Leu Glu Leu  
 305 310 315 320

Tyr Arg Tyr Asn Gly Thr Glu Ile Lys Ile Met Asp Ile Glu Thr Ser  
 325 330 335  
 Asp His Asp Thr Tyr Thr Leu Thr Ser Tyr Pro Asn His Lys Glu Ala  
 340 345 350  
 Leu Leu Leu Leu Thr Asn His Ser Tyr Glu Glu Val Glu Glu Ile Thr  
 355 360 365  
 Lys Ile Pro Lys His Thr Leu Ile Lys Leu Lys Lys His Tyr Phe Lys  
 370 375 380  
 Lys  
 385

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 834 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGACTATATG CAGCAACTTA TTAAAGTTTA GATGATTCAG GTGTTAGTTT AATGAATAAA	60
AATGATGATG ATATTGATGA TTATAACTTA AAATGGTTTT TATTTCTTAT TGATGATGAT	120
CAATATATTA TTACAAGCTA TGCAGCAAAT AATTGTAAAG TTTGGAATGT TAATAATGAT	180
AAAATAAATG TTTCGACTTA TTCTTCAACA AATTCAATAC AAAAATGGCA AATAAAAGCT	240
AATGGTTCTT CATATGTAAT ACAAAGTGAT AATGGAAAAG TCTTAACAGC AGGAACCGGT	300
CAAGCTCTTG GATTGATACG TTAACTGAT GAATCCTCAA ATAATCCCAA TCAACAATGG	360
AATTTAACTT CTGTACAAAC AATTCAACTT CCACAAAAAC CTATAATAGA TACAAAATTA	420
AAAGATTATC CCAAATATTC ACCAACTGGA AATATAGATA ATGGAACATC TCCTCAATTA	480
ATGGGATGGA CATTAGTACC TTGTATTATG GTAAATGATC CAAATATAGA TAAAAATACT	540
CAAATTAAAA CTACTCCATA TTATATTTTA AAAAAATATC AATATTGGCA ACGAGCAGTA	600
GGAAGTAATG TAGCTTTACG TCCACATGAA AAAAAATCAT ATACTTATGA ATGGGGCACA	660
GAAATAGATC AAAAAACAAC AATTATAAAT ACATTAGGAT TTCAAATCAA TATAGATTCA	720
GGAATGAAAT TTGATATACC AGAAGTAGGT GGAGGTACAG ATGAAATAAA AACACAAC TA	780
AATGAAGAAT TAAAAATAGA ATATAGTCAT GAAACTAAAA TAATGGAAAA ATAT	834

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 278 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly	Leu	Tyr	Ala	Ala	Thr	Tyr	Leu	Ser	Leu	Asp	Asp	Ser	Gly	Val	Ser	1	5	10	15
Leu	Met	Asn	Lys	Asn	Asp	Asp	Asp	Ile	Asp	Asp	Tyr	Asn	Leu	Lys	Trp	20	25	30	
Phe	Leu	Phe	Pro	Ile	Asp	Asp	Asp	Gln	Tyr	Ile	Ile	Thr	Ser	Tyr	Ala	35	40	45	
Ala	Asn	Asn	Cys	Lys	Val	Trp	Asn	Val	Asn	Asn	Asp	Lys	Ile	Asn	Val	50	55	60	
Ser	Thr	Tyr	Ser	Ser	Thr	Asn	Ser	Ile	Gln	Lys	Trp	Gln	Ile	Lys	Ala	65	70	75	80
Asn	Gly	Ser	Ser	Tyr	Val	Ile	Gln	Ser	Asp	Asn	Gly	Lys	Val	Leu	Thr	85	90	95	
Ala	Gly	Thr	Gly	Gln	Ala	Leu	Gly	Leu	Ile	Arg	Leu	Thr	Asp	Glu	Ser	100	105	110	
Ser	Asn	Asn	Pro	Asn	Gln	Gln	Trp	Asn	Leu	Thr	Ser	Val	Gln	Thr	Ile	115	120	125	
Gln	Leu	Pro	Gln	Lys	Pro	Ile	Ile	Asp	Thr	Lys	Leu	Lys	Asp	Tyr	Pro	130	135	140	
Lys	Tyr	Ser	Pro	Thr	Gly	Asn	Ile	Asp	Asn	Gly	Thr	Ser	Pro	Gln	Leu	145	150	155	160
Met	Gly	Trp	Thr	Leu	Val	Pro	Cys	Ile	Met	Val	Asn	Asp	Pro	Asn	Ile	165	170	175	
Asp	Lys	Asn	Thr	Gln	Ile	Lys	Thr	Thr	Pro	Tyr	Tyr	Ile	Leu	Lys	Lys	180	185	190	
Tyr	Gln	Tyr	Trp	Gln	Arg	Ala	Val	Gly	Ser	Asn	Val	Ala	Leu	Arg	Pro	195	200	205	
His	Glu	Lys	Lys	Ser	Tyr	Thr	Tyr	Glu	Trp	Gly	Thr	Glu	Ile	Asp	Gln	210	215	220	
Lys	Thr	Thr	Ile	Ile	Asn	Thr	Leu	Gly	Phe	Gln	Ile	Asn	Ile	Asp	Ser	225	230	235	240

Gly Met Lys Phe Asp Ile Pro Glu Val Gly Gly Gly Thr Asp Glu Ile  
 245 250 255

Lys Thr Gln Leu Asn Glu Glu Leu Lys Ile Glu Tyr Ser His Glu Thr  
 260 265 270

Lys Ile Met Glu Lys Tyr  
 275

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 829 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ACATGCAGCA ACTTATTAA GTTTAGATGA TTCAGGTGTT AGTTTAATGA ATAAAAATGA 60

TGATGATATT GATGACTATA ATTTAAGGTG GTTTTATTT CCTATTGATG ATAATCAATA 120

TATTATTACA AGCTACGCAG CGAATAATTG TAAGGTTTGG AATGTTAATA ATGATAAAAT 180

AAATGTTTCA ACTTATTCCT CAACAACTC GATACAGAAA TGGCAATAA AAGCTAATGC 240

TCTTCGTAT GTAATACAAA GTAATAATGG GAAAGTTCTA ACAGCAGGAA CCGGTCATC 300

TCTTGGATTA ATACGTTTAA CGGATGAATC ACCAGATAAT CCCAATCAAC AATGGAATTT 360

AACTCCTGTA CAAACAATTC AACTCCCACC AAAACCTACA ATAGATACAA AGTTAAAAGA 420

TTACCCCAAA TATTCACAAA CTGGCAATAT AGACAAGGGA ACACCTCCTC AATTAATGGG 480

ATGGACATTA ATACCTTGTA TTATGGTAAA TGATCCCAAT ATAGATAAAA ACACTCAAAT 540

CAAACTACT CCATATTATA TTTTAAAAA ATATCAATAT TGGCAACAAG CAGTAGGAAG 600

TAATGTAGCT TTACGTCCGC ATGAAAAAAA ATCATATGCT TATGAGTGGG GTACAGAAAT 660

AGATCAAAA ACACTATCA TTAATACATT AGGATTTTCTG ATTAATATAG ATTCGGGAAT 720

GAAATTTGAT ATACCAGAAG TAGGTGGAGG TACAGATGAA ATAAAAACAC AATTAACGA 780

AGAATTAATA ATAGAATATA GCCGTGAAAC CAAAATAATG GAAAAATAT 829

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 276 amino acids

- (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

His	Ala	Ala	Thr	Tyr	Leu	Ser	Leu	Asp	Asp	Ser	Gly	Val	Ser	Leu	Met
1					5					10				15	
Asn	Lys	Asn	Asp	Asp	Asp	Ile	Asp	Asp	Tyr	Asn	Leu	Arg	Trp	Phe	Leu
		20					25						30		
Phe	Pro	Ile	Asp	Asp	Asn	Gln	Tyr	Ile	Ile	Thr	Ser	Tyr	Ala	Ala	Asn
		35					40					45			
Asn	Cys	Lys	Val	Trp	Asn	Val	Asn	Asn	Asp	Lys	Ile	Asn	Val	Ser	Thr
	50					55					60				
Tyr	Ser	Ser	Thr	Asn	Ser	Ile	Gln	Lys	Trp	Gln	Ile	Lys	Ala	Asn	Ala
65					70					75				80	
Ser	Ser	Tyr	Val	Ile	Gln	Ser	Asn	Asn	Gly	Lys	Val	Leu	Thr	Ala	Gly
			85						90					95	
Thr	Gly	Gln	Ser	Leu	Gly	Leu	Ile	Arg	Leu	Thr	Asp	Glu	Ser	Pro	Asp
			100					105					110		
Asn	Pro	Asn	Gln	Gln	Trp	Asn	Leu	Thr	Pro	Val	Gln	Thr	Ile	Gln	Leu
		115					120					125			
Pro	Pro	Lys	Pro	Thr	Ile	Asp	Thr	Lys	Leu	Lys	Asp	Tyr	Pro	Lys	Tyr
	130					135					140				
Ser	Gln	Thr	Gly	Asn	Ile	Asp	Lys	Gly	Thr	Pro	Pro	Gln	Leu	Met	Gly
145					150					155					160
Trp	Thr	Leu	Ile	Pro	Cys	Ile	Met	Val	Asn	Asp	Pro	Asn	Ile	Asp	Lys
			165						170					175	
Asn	Thr	Gln	Ile	Lys	Thr	Thr	Pro	Tyr	Tyr	Ile	Leu	Lys	Lys	Tyr	Gln
		180						185					190		
Tyr	Trp	Gln	Gln	Ala	Val	Gly	Ser	Asn	Val	Ala	Leu	Arg	Pro	His	Glu
		195					200					205			
Lys	Lys	Ser	Tyr	Ala	Tyr	Glu	Trp	Gly	Thr	Glu	Ile	Asp	Gln	Lys	Thr
	210					215				220					
Thr	Ile	Ile	Asn	Thr	Leu	Gly	Phe	Gln	Ile	Asn	Ile	Asp	Ser	Gly	Met
225				230						235				240	
Lys	Phe	Asp	Ile	Pro	Glu	Val	Gly	Gly	Gly	Thr	Asp	Glu	Ile	Lys	Thr
			245						250					255	

Gln Leu Asn Glu Glu Leu Lys Ile Glu Tyr Ser Arg Glu Thr Lys Ile  
260 265 270

Met Glu Lys Tyr  
275

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Asp Ile Asp Asp Tyr Asn Leu  
1 5

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Trp Phe Leu Phe Pro Ile Asp  
1 5

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 8 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Gln Ile Lys Thr Thr Pro Tyr Tyr  
1 5

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Tyr Glu Trp Gly Thr Glu  
1 5

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GATATNGATG ANTAYAAAYTT N

21

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TGGTTTTTINT TTCNATNGA N

21

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CAAAATNAAAA CNACNCCATA TTAT

24

## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TANGANTGGG GNACAGAA

18

## (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

ATAAATATGGN GTNGITTTNA TTG

24

## (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TTCTGTNCCC CANTCNTA

18

## (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CTCAAAGCGG ATCAGGAG

18

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GCGTATTCGG ATATGCTTGG

20

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 386 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1             5             10             15

Xaa Xaa Xaa Xaa Thr Tyr Leu Ser Leu Asp Asp Ser Gly Val Ser Leu
 20             25             30

Met Xaa Lys Xaa Asp Xaa Asp Ile Asp Asp Tyr Asn Leu Xaa Trp Phe
 35             40             45

Leu Phe Pro Ile Asp Xaa Xaa Gln Tyr Ile Ile Thr Ser Tyr Xaa Ala
 50             55             60

Asn Asn Cys Lys Val Trp Asn Val Xaa Asn Asp Lys Ile Asn Val Ser
 65             70             75             80

Thr Tyr Ser Ser Thr Asn Ser Xaa Gln Lys Trp Gln Ile Lys Ala Xaa
 85             90             95

Xaa Ser Ser Tyr Xaa Ile Gln Ser Xaa Asn Gly Lys Val Leu Thr Ala
 100            105            110

Gly Xaa Gly Gln Xaa Leu Gly Xaa Xaa Arg Leu Thr Asp Glu Xaa Xaa
 115            120            125

```

```

Xaa Asn Xaa Asn Gln Gln Trp Asn Leu Thr Xaa Val Gln Thr Ile Gln
130                      135                      140

Leu Pro Xaa Lys Pro Xaa Ile Asp Xaa Lys Leu Lys Asp Xaa Pro Xaa
145                      150                      155                      160

Tyr Ser Xaa Thr Gly Asn Ile Xaa Xaa Xaa Thr Xaa Pro Gln Leu Met
                      165                      170                      175

Gly Trp Thr Leu Xaa Pro Cys Ile Met Val Asn Asp Xaa Xaa Ile Asp
                      180                      185                      190

Lys Asn Thr Gln Ile Lys Thr Thr Pro Tyr Tyr Ile Xaa Lys Lys Tyr
                      195                      200                      205

Xaa Tyr Trp Xaa Xaa Ala Xaa Gly Ser Asn Val Xaa Leu Xaa Pro His
210                      215                      220

Xaa Lys Xaa Ser Tyr Xaa Tyr Glu Trp Gly Thr Glu Xaa Xaa Gln Lys
225                      230                      235                      240

Thr Thr Ile Ile Asn Thr Xaa Gly Xaa Gln Ile Asn Ile Asp Ser Gly
                      245                      250                      255

Met Lys Phe Xaa Xaa Pro Glu Val Gly Gly Gly Thr Xaa Xaa Ile Lys
260                      265                      270

Thr Gln Leu Xaa Glu Glu Leu Lys Xaa Glu Tyr Ser Xaa Glu Thr Lys
275                      280                      285

Ile Met Xaa Lys Tyr Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
290                      295                      300

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
305                      310                      315                      320

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
325                      330                      335

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
340                      345                      350

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
355                      360                      365

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
370                      375                      380

Xaa Xaa
385

```

## (2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (synthetic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

GNGAAGTNCA TATNGAAATN AATAATAC

28

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 2015 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ATTAATTTTA TGGAGGTTGA TATTTATGTC AGCTCGCGAA GTACACATTG AAATAAACAA	60
TAAAACACGT CATACATTAC AATTAGAGGA TAAAACTAAA CTTAGCGGGC GTAGATGGCG	120
AACATCACCT ACAAATGTTG CTCGTGATAC AATTAAACAA TTTGTAGCAG AATCACATGG	180
TTTTATGACA GGAGTAGAAG GTATTATATA TTTTAGTGTA AACGGAGACG CAGAAATTAG	240
TTTACATTIT GACAATCCTT ATATAGGTTT TAATAAATGT GATGGTCTT CTGATAAACC	300
TGAATATGAA GTTATTACTC AAAGCGGATC AGGAGATAAA TCTCATGTGA CATATACTAT	360
TCAGACAGTA TCTTTACGAT TATAAGGAAA ATTTATAAAA ACTGTATTTT TTTACTAAAT	420
ACCAAAAAAT ACATATTTAT TTTTGGTAT TTTCTAATAT GAAATATGAA TTATAAAAAAT	480
ATTAATAAAA AAGGTGATAA AAATTATGTT AGATACTAAT AAAGTTTATG AAATAAGCAA	540
TCTTGCTAAT GGATTATATA CATCAACTTA TTTAAGTCTT GATGATTGAG GTGTGTAGTTT	600
AATGAGTAAA AAGGATGAAG ATATTGATGA TTACAATTTA AAATGGITTT TATTTCTCTAT	660
TGATAATAAT CAATATATTA TTACAAGCTA TGGAGCTAAT AATTGTAAAG TTTGGAATGT	720
TAAAAATGAT AAAATAAATG TTTCAACTTA TTCTTCAACA AACTCTGTAC AAAAAATGCA	780
AATAAAGCT AAAGATTCTT CATATATAAT ACAAAGTGAT AATGGAAAGG TCTTAACAGC	840
AGGAGTAGGT CAATCTCTTG GAATAGTACG CCTAAGTGAT GAATTTCCAG AGAATCTTAA	900
CCAACAATGG AATTTAACTC CTGTACAAAC AATTCAACTC CCACAAAAAC CTAATAAGTA	960
TGAAAAATTA AAAGATCATC CTGAATATTC AGAAACCGGA AATATAAATC CTAATAACAC	1020
TCCTCAATTA ATGGGATGGA CATTAGTACC TTGTATTATG GTAAATGATT CAAAAATAGA	1080

TAAAAAACT CAAATTAATA CTACTCCATA TTATATTTTT AAAAAATATA AATACTGGAA	1140
TCTAGCAAAA GGAAGTAATG TATCTTTACT TCCACATCAA AAAAGATCAT ATGATTATGA	1200
ATGGGGTACA GAAAAAATC AAAAAACAAC TATTATTAAT ACAGTAGGAT TGCAAAATTAA	1260
TATAGATTCA GGAATGAAAT TTGAAGTACC AGAAGTAGGA GGAGGTACAG AAGACATAAA	1320
AACACAATTA ACTGAAGAAT TAAAAGTTGA ATATAGCACT GAAACCAAAA TAATGACGAA	1380
ATATCAAGAA CACTCAGAGA TAGATAATCC AACTAATCAA CCAATGAATT CTATAGGACT	1440
TCTTATTTAT ACTTCTTTAG AATTATATCG ATATAACGGT ACAGAAATTA AGATAATGGA	1500
CATAGAAACT TCAGATCATG ATACTTACAC TCTTACTTCT TATCCAAATC ATAAAGAAGC	1560
ATTATTACTT CTCACAAACC ATTCGTATGA AGAAGTAGAA GAAATAACAA AAATACCTAA	1620
GCATACACTT ATAAAAATGA AAAAACATTA TTTTAAAAAA TAAAAAACAT AATATATAAA	1680
TGACTGATTA ATATCTCTCG AAAAGGTTCT GGTGCAAAAA TAGTGGGATA TGAAAAAAGC	1740
AAAAGATTCC TAACGGAATG GAACATTAGG CTGTTAAATC AAAAAGTTTA TTGATAAAAT	1800
ATATCTGCCT TTGGACAGAC TTCGCCCTT GGAGAGTTTG TCCTTTTTTG ACCATATGCA	1860
TAGCTTCTAT TCCGGCAATC ATTTTGTAG CTGTTTGCAA GGATTTTAAT CCAAGCATAT	1920
CCGAATACGC TTTTGTATAA CCGATGTCTT GTTCAATGAT ATTGTTTAAT ATTTTCACAC	1980
GAATTGGCTA CTGTGCGGTA TCCTGTCTCC TTTAT	2015

## (2) INFORMATION FOR SEQ ID NO:31:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 360 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATGTCAGCTC GCGAAGTACA CATTGAAATA AACAATAAAA CACGTCATAC ATTACAATTA	60
GAGGATAAAA CTAAACTTAG CGGCGGTAGA TGGCGAACAT CACCTACAAA TGTGTCTCGT	120
GATACAATTA AAACATTTGT AGCAGAATCA CATGGTTTTA TGACAGGAGT AGAAGGTATT	180
ATATATTTTA GTGTAAACGG AGACGCAGAA ATTAGTTTAC ATTTTGACAA TCCTTATATA	240
GGTTCTAATA AATGTGATGG TTCTTCTGAT AAACCTGAAT ATGAAGTTAT TACTCAAAGC	300
GGATCAGGAG ATAAATCTCA TGTGACATAT ACTATTGAGA CAGTATCTTT ACGATTATAA	360

## (2) INFORMATION FOR SEQ ID NO:32:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 119 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

```

Met Ser Ala Arg Glu Val His Ile Glu Ile Asn Asn Lys Thr Arg His
 1             5             10             15

Thr Leu Gln Leu Glu Asp Lys Thr Lys Leu Ser Gly Gly Arg Trp Arg
      20             25             30

Thr Ser Pro Thr Asn Val Ala Arg Asp Thr Ile Lys Thr Phe Val Ala
      35             40             45

Glu Ser His Gly Phe Met Thr Gly Val Glu Gly Ile Ile Tyr Phe Ser
      50             55             60

Val Asn Gly Asp Ala Glu Ile Ser Leu His Phe Asp Asn Pro Tyr Ile
      65             70             75             80

Gly Ser Asn Lys Cys Asp Gly Ser Ser Asp Lys Pro Glu Tyr Glu Val
      85             90             95

Ile Thr Gln Ser Gly Ser Gly Asp Lys Ser His Val Thr Tyr Thr Ile
      100            105            110

Gln Thr Val Ser Leu Arg Leu
      115

```

## (2) INFORMATION FOR SEQ ID NO:33:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CATGAGATTT ATCTCCTGAT CCGC

24

## (2) INFORMATION FOR SEQ ID NO:34:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2230 base pairs
- (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

ACTATGACAA TGATTATGAC TGCTGATGAA TTAGCTTTAT CAATACCAGG ATATTCTAAA	60
CCATCAAATA TAACAGGAGA TAAAGTAAA CATACTTAT TTAATAATAT AATTGGAGAT	120
ATTCAAATAA AAGATCAAGC AACATTTGGG GTTGTTTTGG ATCCCCCTCT TAATCGTATT	180
TCAGGGGCTG AAGAATCAAG TAAGTTTATT GATGTATATT ATCCTTCTGA AGATAGTAAC	240
CTTAAATATT ATCAATTAT AAAAGTAGCA ATTGATTTTG ATATTAATGA AGATTTTATT	300
AATTTTAATA ATCATGACAA TATAGGGATA TTTAATTTTG TTACACGAAA TTTTTTATTA	360
AATAATGAAA ATGATTAATA AAAAATTTAA TTTGTATAAT ATGTTTATTT TTTGAAAATT	420
GAATGCATAT ATTAATCGAG TATGTGTAAT AAATTTTAAT TTTATGGAGG TTGATATTTA	480
TGTCAGCACG TGAAGTACAC ATTGATGTAA ATAATAAGAC AGGTCATACA TTACAATTAG	540
AAGATAAAAC AAAACTTGAT GGTGGTAGAT GSCGAACATC ACCTACAAAT GTTGCTAATG	600
ATCAAAATTA AACATTTGTA GCAGAAATCAC ATGGTTTTAT GACAGGTACA GAAGGTAATA	660
TATATTATAG TATAAATGGA GAAGCAGAAA TTAGTTTATA TTTTGACAAAT CCTTATTCAG	720
GTTCTAATAA ATATGATGGG CATTCCAATA AAAATCAATA TGAAGTTATT ACCCAAGGAG	780
GATCAGGAAA TCAATCTCAT GTTACGTATA CTATTCAAAC TGATATCTCA CGATATGGGA	840
ATAATTCATA AAAAAATATT TTTTTTACG AAAATACCAA AAAAAATTTT TTGGTATTTT	900
CTAATATAAT TCATAAATAT TTTAATAATA AAATTATAAG AAAAGGTGAT AAATATTATG	960
TTAGATACTA ATAAAAATTA TGAAATAAGT AATTATGCTA ATGGATTACA TGCAGCAACT	1020
TATTTAAGTT TAGATGATTC AGGTGTTAGT TTAATGAATA AAAATGATGA TGATATTGAT	1080
GACTATAATT TAAGGTGGTT TTTATTTCTT ATTGATGATA ATCAATATAT TATTACAAGC	1140
TACGCAGCGA ATAATTGTAA GGTTTGGAAT GTTAATAATG ATAAAATAAA TGTTCACACT	1200
TATTCCTCAA CAACTCGAT ACAGAAATGG CAAATAAAAG CTAATGCTTC TTCGTATGTA	1260
ATACAAAGTA ATAATGGGAA AGTTCTAACA GCAGGAACCG GTCAATCTCT TGGATTATAA	1320
CGTTTAACGG ATGAATCACC AGATAATCCC AATCAACAAT GGAATTTAAC TCCTGTACAA	1380
ACAATTCAAC TCCCACCAA ACCTACAATA GATACAAAGT TAAAAGATTA CCCCAAATAT	1440
TCACAAACTG GCAATATAGA CAAGGAACA CCTCCTCAAT TAATGGGATG GACATTAATA	1500

CCTTGTTATTA TGGTAAATGA TCCAAATATA GATAAAAACA CTCAAATCAA AACTACTCCA	1560
TATTATATTT TAAAAAATA TCAATATTGG CAACAAGCAG TAGGAAGTAA TGTAGCTTTA	1620
CGTCCGCATG AAAAAAATC ATATGCTTAT GAGTGGGGTA CAGAAATAGA TCAAAAAACA	1680
ACTATCATTAT ATACATTAGG ATTTGAGATT AATATAGATT CGGGAATGAA ATTTGATATA	1740
CCAGAAGTAG GTGGAGGTAC AGATGAAATA AAAACACAAT TAAACGAAGA ATTAATAATA	1800
GAATATAGCC GTGAAACCAA AATAATGGAA AAATATCAGG ACAAATCAGA GATAGATAAT	1860
CCAAGTATC AATCAATGAA TTCTATAGGA TTCCTCACTA TTACTTCTTT AGAATTATAT	1920
CGATATAATG GTTCGGAAT TAGTGTAATG AAAATTCAAA CTTCAGATAA TGATACTTAC	1980
AATGTGACCT CTTATCCAGA TCATCAACAA GCTCTATTAC TTCTTACAAA TCATTATAT	2040
GAAGAAGTAG AAGAAATAAC AAATATCCC AAAATATCAC TGAAAAAATT AAAAAAATAT	2100
TATTTTTTAA ACATAATTAT ATTTTGATAG CTTTTTAAAA ATAAAGATTG TTCAAAGTAA	2160
AATGAAAGAA AATCTTTTAT GAAACTTTAA TACAATAAAA GAGGAATATT TTCTTATAAG	2220
TACTTCCTTG	2230

## (2) INFORMATION FOR SEQ ID NO:35:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

ATGTCAGCAC GTGAAGTACA CATTGATGTA AATAATAAGA CAGGTCATAC ATTACAATTA	60
GAAGATAAAA CAAAACCTGA TGGTGGTAGA TGGCGAATAC CACCTACAAA TGTGCTAAT	120
GATCAAATTA AAACATTTGT AGCAGAATCA CATGGTTTTA TGACAGGTAC AGAAGGTACT	180
ATATATTATA GTATAAATGG AGAAGCAGAA ATTAGTTTAT ATTTTGACAA TCCTTATTCA	240
GGTTCTAATA AATATGATGG GCATTCCAAT AAAAATCAAT ATGAAGTTAT TACCCAAGGA	300
GGATCAGGAA ATCAATCTCA TGTTACGTAT ACTATTCAAA CTGTATCTTC ACGATATGGG	360
AATAATTCAT AA	372

## (2) INFORMATION FOR SEQ ID NO:36:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 123 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

```

Met Ser Ala Arg Glu Val His Ile Asp Val Asn Asn Lys Thr Gly His
 1             5             10             15

Thr Leu Gln Leu Glu Asp Lys Thr Lys Leu Asp Gly Gly Arg Trp Arg
          20             25             30

Thr Ser Pro Thr Asn Val Ala Asn Asp Gln Ile Lys Thr Phe Val Ala
          35             40             45

Glu Ser His Gly Phe Met Thr Gly Thr Glu Gly Thr Ile Tyr Tyr Ser
 50             55             60

Ile Asn Gly Glu Ala Glu Ile Ser Leu Tyr Phe Asp Asn Pro Tyr Ser
 65             70             75             80

Gly Ser Asn Lys Tyr Asp Gly His Ser Asn Lys Asn Gln Tyr Glu Val
          85             90             95

Ile Thr Gln Gly Gly Ser Gly Asn Gln Ser His Val Thr Tyr Thr Ile
          100            105            110

Gln Thr Val Ser Ser Arg Tyr Gly Asn Asn Ser
          115            120

```

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1152 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

```

ATGTTAGATA CTAATAAAAT TTATGAAATA AGTAATTATG CTAATGGATT ACATGCAGCA      60
ACTTATTITAA GTTTAGATGA TTCAGGTGTT AGTTTAATGA ATAAAAATGA TGATGATATT      120
GATGACTATA ATTTAAGGTG GTTTTTATTT CCTATTGATG ATAATCAATA TATTATTACA      180
AGCTACGCAG CGAATAATTG TAAGGTTTGG AATGTTAATA ATGATAAAAT AAATGTTTCA      240
ACTTATTCTT CAACAAACTC GATACAGAAA TGGCAAATAA AAGCTAATGC TTCTTCGTAT      300
GTAATACAAA GTAATAATGG GAAAGTTCTA ACAGCAGGAA CCGGTCAATC TCTTGGATTA      360

```

ATACGTTTAA	CGGATGAATC	ACCAGATAAT	CCCAATCAAC	AATGGAATTT	AACTCCTGTA	420
CAAACAATT	CACTCCACC	AAAACCTACA	ATAGATACAA	AGTTAAAAGA	TTACCCCAAA	480
TATTCACAAA	CTGGCAATAT	AGACAAGGGA	ACACCTCCTC	AATTAATGGG	ATGGACATTA	540
ATACCTTGTA	TTATGGTAAA	TGATCCAAAT	ATAGATAAAA	ACACTCAAAT	CAAAACTACT	600
CCATATTATA	TTTTAAAAAA	ATATCAATAT	TGGCAACAAG	CAGTAGGAAG	TAATGTAGCT	660
TTACGTCCGC	ATGAAAAAAA	ATCATATGCT	TATGAGTGGG	GTACAGAAAT	AGATCAAAAA	720
ACAACTATCA	TTAATACATT	AGGATTTCAG	ATTAATATAG	ATTCGGGAAT	GAAATTTGAT	780
ATACCAGAAG	TAGGTGGAGG	TACAGATGAA	ATAAAAACAC	AATTAACAGA	AGAATTAAAA	840
ATAGAATATA	GCCGTGAAAC	CAAAATAATG	GAAAAATATC	AGGAACAATC	AGAGATAGAT	900
AATCCAACCTG	ATCAATCAAT	GAATTCTATA	GGATTCTCTA	CTATTACTTC	TTTAGAATTA	960
TATCGATATA	ATGGTTCGGA	AATTAGTGTA	ATGAAAATTC	AAACTTCAGA	TAATGATACT	1020
TACAATGTGA	CCTCTTATCC	AGATCATCAA	CAAGCTCTAT	TACTTCTTAC	AAATCATTTCA	1080
TATGAAGAAG	TAGAAGAAAT	AACAAATATT	CCCAAAATAT	CAGTAAAAAA	ATTAAAAAAA	1140
TATTATTTTT	AA					1155

(2) INFORMATION FOR SEO ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEO ID NO:38:

Met	Leu	Asp	Thr	Asn	Lys	Ile	Tyr	Glu	Ile	Ser	Asn	Tyr	Ala	Asn	Gly
1				5					10					15	
Leu	His	Ala	Ala	Thr	Tyr	Leu	Ser	Leu	Asp	Asp	Ser	Gly	Val	Ser	Leu
		20						25					30		
Met	Asn	Lys	Asn	Asp	Asp	Asp	Ile	Asp	Asp	Tyr	Asn	Leu	Arg	Trp	Phe
		35					40					45			
Leu	Phe	Pro	Ile	Asp	Asp	Asn	Gln	Tyr	Ile	Ile	Thr	Ser	Tyr	Ala	Ala
	50					55					60				
Asn	Asn	Cys	Lys	Val	Trp	Asn	Val	Asn	Asn	Asp	Lys	Ile	Asn	Val	Ser
65					70					75				80	

Thr Tyr Ser Ser Thr Asn Ser Ile Gln Lys Trp Gln Ile Lys Ala Asn  
 85 90 95  
 Ala Ser Ser Tyr Val Ile Gln Ser Asn Asn Gly Lys Val Leu Thr Ala  
 100 105 110  
 Gly Thr Gly Gln Ser Leu Gly Leu Ile Arg Leu Thr Asp Glu Ser Pro  
 115 120 125  
 Asp Asn Pro Asn Gln Gln Trp Asn Leu Thr Pro Val Gln Thr Ile Gln  
 130 135 140  
 Leu Pro Pro Lys Pro Thr Ile Asp Thr Lys Leu Lys Asp Tyr Pro Lys  
 145 150 155 160  
 Tyr Ser Gln Thr Gly Asn Ile Asp Lys Gly Thr Pro Pro Gln Leu Met  
 165 170 175  
 Gly Trp Thr Leu Ile Pro Cys Ile Met Val Asn Asp Pro Asn Ile Asp  
 180 185 190  
 Lys Asn Thr Gln Ile Lys Thr Thr Pro Tyr Tyr Ile Leu Lys Lys Tyr  
 195 200 205  
 Gln Tyr Trp Gln Gln Ala Val Gly Ser Asn Val Ala Leu Arg Pro His  
 210 215 220  
 Glu Lys Lys Ser Tyr Ala Tyr Glu Trp Gly Thr Glu Ile Asp Gln Lys  
 225 230 235 240  
 Thr Thr Ile Ile Asn Thr Leu Gly Phe Gln Ile Asn Ile Asp Ser Gly  
 245 250 255  
 Met Lys Phe Asp Ile Pro Glu Val Gly Gly Gly Thr Asp Glu Ile Lys  
 260 265 270  
 Thr Gln Leu Asn Glu Glu Leu Lys Ile Glu Tyr Ser Arg Glu Thr Lys  
 275 280 285  
 Ile Met Glu Lys Tyr Gln Glu Gln Ser Glu Ile Asp Asn Pro Thr Asp  
 290 295 300  
 Gln Ser Met Asn Ser Ile Gly Phe Leu Thr Ile Thr Ser Leu Glu Leu  
 305 310 315 320  
 Tyr Arg Tyr Asn Gly Ser Glu Ile Ser Val Met Lys Ile Gln Thr Ser  
 325 330 335  
 Asp Asn Asp Thr Tyr Asn Val Thr Ser Tyr Pro Asp His Gln Gln Ala  
 340 345 350  
 Leu Leu Leu Leu Thr Asn His Ser Tyr Glu Glu Val Glu Glu Ile Thr  
 355 360 365  
 Asn Ile Pro Lys Ile Ser Leu Lys Lys Leu Lys Lys Tyr Tyr Phe  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:39:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2132 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

GTATTTTCAGG GGGTGAAGAT TCAAGTAAGT TTATTGATGT ATATTATCCT TTTGAAGATA	60
GTAATTTTAA ATATTATCAA TTTATAAAAG TAGCAATTGA TTTTGATATT AATGAAGATT	120
TTATTAAATTT TAATAATCAT GACAATATAG GGATATTTAA TTTTGTGTACA CGAAATTTTT	180
TATTAATAAA TGAAAATGAT GAATAAAAAA TTAAATTGTG TTATTATGTT TATTTTTTGA	240
AAATTGAATG CATATATTAA TCGAGTAGTG ATAATAAATT TTAATTTTAT GGAGGTTGAT	300
ATTTATGTCA GCACGTGAAG TACACATTGA TGTAAATAAT AAGACAGGTC ATACATTACA	360
ATTAGAAGAT AAAACAAAAC TTGATGGTGG TAGATGGCGA ACATCACCTA CAAATGTTGC	420
TAATGATCAA ATTAAACAT TTGTAGCAGA ATCAAATGGT TTTATGACAG GTACAGAAGG	480
TACTATATAT TATAGTATAA ATGGAGAAGC AGAAATTAGT TTATATTTTG ACAATCCTTT	540
TGCAGGTCTT AATAAATATG ATGGACATTC CAATAAATCT CAATATGAAA TTATTACCCA	600
AGGAGGATCA GGAAATCAAT CTCATGTTAC GTATACTATT CAAACCACAT CCTCACGATA	660
TGGGCATAAA TCATAACAAA TAATTTTTTA CGAAAATACC AAAAAATAAA TATTTTTTGG	720
TATTTTCTAA TATAAATTAC AAATATATTA ATAATAAAAT TATAAGAAAA GGTGATAAAG	780
ATTATGTTAG ATACTAATAA AGTTTATGAA ATAAGCAATC ATGCTAATGG ACTATATGCA	840
GCAACTTATT TAAGTTTAGA TGATTCAAGT GTTAGTTTAA TGAATAAAAA TGATGATGAT	900
ATTGATGATT ATAACTTAAA ATGGTTTTTA TTTCCTATTG ATGATGATCA ATATATTATT	960
ACAAGCTATG CAGCAAAATA TTGTAAAGTT TGGAAATGTTA ATAATGATAA AATAAATGTT	1020
TCGACTTATT CTTCACAAA TTCAATACAA AAATGGCAAA TAAAAGCTAA TGGTCTTCA	1080
TATGTAATAC AAAGTGATAA TGGAAAAGTC TTAACAGCAG GAACCGGTCA AGCTCTTGGA	1140
TTGATACGTT TAACTGATGA ATCCTCAAAT AATCCCAATC AACAATGGAA TTAACTTCT	1200
GTACAAACAA TTCAACTTCC ACAAAAAACCT ATAATAGATA CAAATTTAAA AGATTATCCC	1260
AAATATTCAC CAACTGGAAA TATAGATAAT GGAACATCTC CTCAATTAAAT GGGATGGACA	1320
TTAGTACCTT GTATTATGGT AAATGATCCA AATATAGATA AAAATACTCA AATTAAAACT	1380

ACTCCATATT ATATTTTAAA AAAATATCAA TATTGGCAAC GAGCAGTAGG AAGTAATGTA	1440
GCTTTACGTC CACATGAAAA AAAATCATAT ACTTATGAAT GGGGCACAGA AATAGATCAA	1500
AAAACAACAA TTATAAATAC ATTAGGATTT CAAATCAATA TAGATTGAGG AATGAAATTT	1560
GATATACCAG AAGTAGGTGG AGGTACAGAT GAAATAAAAA CACAACATAA TGAAGAATTA	1620
AAAATAGAAT ATAGTCATGA AACTAAAATA ATGGAAAAAT ATCAAGAACA ATCTGAAATA	1680
GATAATCCAA CTGATCAATC AATGAATTCT ATAGGATTTT TTACTATTAC TTCTTAGAA	1740
TTATATAGAT ATAATGGCTC AGAAATTCGT ATAATGCAAA TTCAAACCTC AGATAATGAT	1800
ACTTATAATG TTACTTCTTA TCCAAATCAT CAACAAGCTT TATTACTTCT TACAAATCAT	1860
TCATATGAAG AAGTAGAAGA AATAACAAAT ATTCCTAAAA GTACACTAAA AAAATTAAAA	1920
AAATATTATT TTTAAATATT GAAATTAGAA ATTATCTAAA ACAAAACGAA AGATAATTTA	1980
ATCTTTAATT ATTTGTAAGA TAATCGTATT TTATTGTAT TAATTTTAT ACAATATAAA	2040
GTAATATCTG TACGTGAAAT TGGTTTCGT TCAATATCTA ATCTCATCTC ATGTATTACA	2100
TGCGTAATAC CTTCTGTTC TGCTTCTACA AG	2132

## (2) INFORMATION FOR SEQ ID NO:40:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

ATGTCAGCAC GTGAAGTACA CATGTATGTA AATAATAAGA CAGGTCATAC ATTACAATTA	60
GAAGATAAAA CAAAACTTGA TGGTGGTAGA TGGCGAACAT CACCTACAAA TGTTGCTAAT	120
GATCAAATTA AAACATTGTG AGCAGAATCA AATGGTTTTA TGACAGGTAC AGAAGGTACT	180
ATATATTATA GTATAAATGG AGAAGCAGAA ATTAGTTTAT ATTTTGACAA TCCTTTTGCA	240
GGTTCTAATA AATATGATGG ACATTCCAAT AAATCTCAAT ATGAAATTAT TACCCAAGGA	300
GGATCAGGAA ATCAATCTCA TGTTACGTAT ACTATTCAAA CCACATCCTC ACGATATGGG	360
CATAAATCAT AA	372

## (2) INFORMATION FOR SEQ ID NO:41:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 123 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

```

Met Ser Ala Arg Glu Val His Ile Asp Val Asn Asn Lys Thr Gly His
 1             5             10             15

Thr Leu Gln Leu Glu Asp Lys Thr Lys Leu Asp Gly Gly Arg Trp Arg
          20             25             30

Thr Ser Pro Thr Asn Val Ala Asn Asp Gln Ile Lys Thr Phe Val Ala
          35             40             45

Glu Ser Asn Gly Phe Met Thr Gly Thr Glu Gly Thr Ile Tyr Tyr Ser
 50             55             60

Ile Asn Gly Glu Ala Glu Ile Ser Leu Tyr Phe Asp Asn Pro Phe Ala
 65             70             75             80

Gly Ser Asn Lys Tyr Asp Gly His Ser Asn Lys Ser Gln Tyr Glu Ile
          85             90             95

Ile Thr Gln Gly Gly Ser Gly Asn Gln Ser His Val Thr Tyr Thr Ile
          100            105            110

Gln Thr Thr Ser Ser Arg Tyr Gly His Lys Ser
          115            120

```

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1152 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

```

ATGTTAGATA CTAATAAAGT TTATGAAATA AGCAATCATG CTAATGGACT ATATGCAGCA      60
ACTTATTTTAA GTTTAGATGA TTCAGGTGTT AGTTTAATGA ATAAAAATGA TGATGATATT      120
GATGATTATA ACTTAAATG GTTTTTATTT CCTATTGATG ATGATCAATA TATTATTACA      180
AGCTATGCAG CAAATAATTG TAAAGTTTGG AATGTTAATA ATGATAAAAT AAATGTTTCG      240
ACTTATTCTT CAACAAATTC AATACAAAAA TGGCAAATAA AAGCTAATGG TTCTTCATAT      300
GTAATACAAA GTGATAATGG AAAAGTCTTA ACAGCAGGAA CCGGTCAAGC TCTTGGATTG      360

```

ATACGTTTAA CTGATGAATC CTCAAATAAT CCCAATCAAC AATGGAATTT AACTTCTGTA	420
CAACAACATTC AACTTCCACA AAAACCTATA ATAGATACAA AATTAAAAGA TTATCCCAA	480
TATTCAACAA CTGGAATAT AGATAATGGA ACATCTCCTC AATTAATGGG ATGGACATTA	540
GTACCTTGTA TTATGGTAAA TGATCCAAAT ATAGATAAAA ATACTCAAAT TAAACTACT	600
CCATATTATA TTTTAAAAA ATATCAATAT TGCAACGAG CAGTAGGAAG TAATGTAGCT	660
TTACGTCCAC ATGAAAAAA ATCATATACT TATGAATGGG GCACAGAAAT AGATCAAAAA	720
ACAACAATTA TAAATACATT AGGATTTCOA ATCAATATAG ATTCAGGAAT GAAATTTGAT	780
ATACCAGAAG TAGGTGGAGG TACAGATGAA ATAAAAACAC AACTAAATGA AGAATTAAAA	840
ATAGAATATA GTCATGAAAC TAAAAAATG GAAAAATATC AAGAACAATC TGAATAGAT	900
AAATCCAACTG ATCAATCAAT GAATTCATATA GGATTTCCTA CTATTACTTC CTTAGAATTA	960
TATAGATATA ATGGCTCAGA AATTCGTATA ATGCAAAATC AACCTCAGA TAATGATACT	1020
TATAATGTGA CTTCTTATCC AAATCATCAA CAAGCTTTAT TACTTCTTAC AAATCATTCA	1080
TATGAAGAAG TAGAAGAAAT AACAAATATT CCTAAAAGTA CACTAAAAAA ATTAAAAAAA	1140
TATTATTTTT AA	1152

## (2) INFORMATION FOR SEQ ID NO:43:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Met	Leu	Asp	Thr	Asn	Lys	Val	Tyr	Glu	Ile	Ser	Asn	His	Ala	Asn	Gly
1				5				10						15	
Leu	Tyr	Ala	Ala	Thr	Tyr	Leu	Ser	Leu	Asp	Asp	Ser	Gly	Val	Ser	Leu
		20					25						30		
Met	Asn	Lys	Asn	Asp	Asp	Asp	Ile	Asp	Asp	Tyr	Asn	Leu	Lys	Trp	Phe
		35					40					45			
Leu	Phe	Pro	Ile	Asp	Asp	Asp	Gln	Tyr	Ile	Ile	Thr	Ser	Tyr	Ala	Ala
	50					55					60				
Asn	Asn	Cys	Lys	Val	Trp	Asn	Val	Asn	Asn	Asp	Lys	Ile	Asn	Val	Ser
65					70					75					80

Thr Tyr Ser Ser Thr Asn Ser Ile Gln Lys Trp Gln Ile Lys Ala Asn  
 85 90 95  
 Gly Ser Ser Tyr Val Ile Gln Ser Asp Asn Gly Lys Val Leu Thr Ala  
 100 105 110  
 Gly Thr Gly Gln Ala Leu Gly Leu Ile Arg Leu Thr Asp Glu Ser Ser  
 115 120 125  
 Asn Asn Pro Asn Gln Gln Trp Asn Leu Thr Ser Val Gln Thr Ile Gln  
 130 135 140  
 Leu Pro Gln Lys Pro Ile Ile Asp Thr Lys Leu Lys Asp Tyr Pro Lys  
 145 150 155 160  
 Tyr Ser Pro Thr Gly Asn Ile Asp Asn Gly Thr Ser Pro Gln Leu Met  
 165 170 175  
 Gly Trp Thr Leu Val Pro Cys Ile Met Val Asn Asp Pro Asn Ile Asp  
 180 185 190  
 Lys Asn Thr Gln Ile Lys Thr Thr Pro Tyr Tyr Ile Leu Lys Lys Tyr  
 195 200 205  
 Gln Tyr Trp Gln Arg Ala Val Gly Ser Asn Val Ala Leu Arg Pro His  
 210 215 220  
 Glu Lys Lys Ser Tyr Thr Tyr Glu Trp Gly Thr Glu Ile Asp Gln Lys  
 225 230 235 240  
 Thr Thr Ile Ile Asn Thr Leu Gly Phe Gln Ile Asn Ile Asp Ser Gly  
 245 250 255  
 Met Lys Phe Asp Ile Pro Glu Val Gly Gly Gly Thr Asp Glu Ile Lys  
 260 265 270  
 Thr Gln Leu Asn Glu Glu Leu Lys Ile Glu Tyr Ser His Glu Thr Lys  
 275 280 285  
 Ile Met Glu Lys Tyr Gln Glu Gln Ser Glu Ile Asp Asn Pro Thr Asp  
 290 295 300  
 Gln Ser Met Asn Ser Ile Gly Phe Leu Thr Ile Thr Ser Leu Glu Leu  
 305 310 315 320  
 Tyr Arg Tyr Asn Gly Ser Glu Ile Arg Ile Met Gln Ile Gln Thr Ser  
 325 330 335  
 Asp Asn Asp Thr Tyr Asn Val Thr Ser Tyr Pro Asn His Gln Gln Ala  
 340 345 350  
 Leu Leu Leu Leu Thr Asn His Ser Tyr Glu Glu Val Glu Glu Ile Thr  
 355 360 365  
 Asn Ile Pro Lys Ser Thr Leu Lys Lys Leu Lys Lys Tyr Tyr Phe  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:44:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 360 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

ATGTCCGCC	CGGAGGTGCA	CATCGAGATC	AACAACAAGA	CCGCCACAC	CCTCCAGCTC	60
GAGGACAAGA	CCAAGCTCTC	CGGCGCAGG	TGGCGCACCT	CCCCGACCAA	CGTGGCCCGC	120
GACACCATCA	AGACGTTCTG	GGCGGAGTCC	CACGGCTTCA	TGACCGGCGT	CGAGGCGATC	180
ATCTACTTCT	CCGTGAACGG	CGACGCCGAG	ATCTCCCTCC	ACTTCGACAA	CCCGTACATC	240
GGCTCCAACA	AGTSCGACGG	CTCTCCGAC	AAGCCCGAGT	ACGAGGTGAT	CACCCAGTCC	300
GGCTCCGGCG	ACAAGTCCCA	CGTGACCTAC	ACCATCCAGA	CCGTGTCCCT	CCGCCTCTGA	360

## (2) INFORMATION FOR SEQ ID NO:45:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1158 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ATGCTCGACA	CCAACAAGGT	GTACGAGATC	TCCAACCTCG	CCAACGGCCT	CTACACCTCC	60
ACCTACCTCT	CCCTCGACGA	CTCCGGCGTG	TCCCTCATGT	CCAAGAAGGA	CGAGGACATC	120
GAGACTACA	ACCTCAAGTG	GTTCTCTTTC	CCGATCGACA	ACAACCGATA	CATCATCACC	180
TCCTACGGCG	CCAACAACCTG	CAAGGTGTGG	AACGTGAAGA	ACGACAAGAT	CAACGTGTCC	240
ACCTACTCCT	CCACCAACTC	CGTGCGAAG	TGGCAGATCA	AGGCCAAGGA	CTCCTCTTAC	300
ATCATCCAGT	CCGACAACGG	CAAGGTGCTC	ACCGCGGGCG	TGGGCCAGTC	CCTCGGCATC	360
GTGCGCTCA	CCGACGAGTT	CCCGGAGAAC	TCCAACCAGC	AATGGAACCT	CACCCCGGTG	420
CAGACCATCC	AGCTCCCGCA	GAAGCCGAAG	ATCGACGAGA	AGCTCAAGGA	CCACCCGGAG	480
TACTCCGAGA	CCGGCAACAT	CAACCCGAAG	ACCACCCCGC	AGCTCATGGG	CTGGACCTTC	540
GTGCCGTGCA	TCATGTTGAA	CGACTCCAAG	ATCGACAAGA	ACACCCAGAT	CAAGACCACC	600

CCGTACTACA TCTTCAAGAA ATACAAGTAC TGGAACCTCG CCAAGGGCTC CAACGTGTCC	660
CTCCTCCCGC ACCAGAAGCG CAGCTACGAC TACGAGTGGG GCACCGAGAA GAACCAGAAG	720
ACCACCATCA TCAACACCGT GGGCCTGCAG ATCAACATCG ACTCGGGGAT GAAGTTCGAG	780
GTGCCGGAGG TGGGCGGCGG CACCGAGGAC ATCAAGACCC AGCTCACCGA GGAGCTGAAG	840
GTGGAGTACT CCACCGAGAC CAAGATCATG ACCAAGTACC AGGAGCACTC CGAGATCGAC	900
AACCCGACCA ACCAGCCGAT GAACTCCATC GGCTCCTCA TCTACACCTC CCTCGAGCTG	960
TACCGCTACA ACGGCACCGA GATCAAGATC ATGGACATCG AGACCTCCGA CCACGACACC	1020
TACACCTCA CCTCTACCC GAACCACAAG GAGGCGCTGC TGCTGCTGAC CAACCACTCC	1080
TACGAGGAGG TGGAGGAGAT CACCAAGATC CCGAAGCACA CCCTCATCAA GCTCAAGAAG	1140
CACTACTTCA AGAAGTGA	1158

Claims

1           1. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2 active against a non-mammalian pest wherein said nucleotide sequence hybridizes under  
3 stringent conditions with a nucleotide sequence selected from the group consisting of: DNA  
4 which encodes SEQ ID NO. 2; DNA which encodes SEQ ID NO. 4; DNA which encodes SEQ  
5 ID NO. 6; SEQ ID NO. 8; SEQ ID NO. 10; DNA which encodes SEQ ID NO. 11; SEQ ID NO.  
6 12; DNA which encodes SEQ ID NO. 13; SEQ ID NO. 14; DNA which encodes SEQ ID NO.  
7 15; DNA which encodes SEQ ID NO. 16; DNA which encodes SEQ ID NO. 17; DNA which  
8 encodes SEQ ID NO. 18; DNA which encodes SEQ ID NO. 19; SEQ ID NO. 20; SEQ ID NO.  
9 21; SEQ ID NO. 22; SEQ ID NO. 23; SEQ ID NO. 24; SEQ ID NO. 25; SEQ ID NO. 26; SEQ  
10 ID NO. 27; DNA which encodes a pesticidal portion of SEQ ID NO. 28; SEQ ID NO. 37; DNA  
11 which encodes SEQ ID NO. 38; SEQ ID NO. 42; and DNA which encodes SEQ ID NO. 43.

1           2. The isolated polynucleotide, according to claim 1, wherein said nucleotide sequence  
2 hybridizes with DNA which encodes SEQ ID NO. 2.

1           3. The isolated polynucleotide, according to claim 1, wherein said nucleotide sequence  
2 hybridizes with SEQ ID NO. 10.

1           4. The isolated polynucleotide, according to claim 1, wherein said toxin has a molecular  
2 weight of approximately 40-50 kDa.

1           5. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2 active against a non-mammalian pest wherein said toxin immunoreacts with an antibody to an  
3 approximately 40-50 kDa toxin from a *Bacillus thuringiensis* isolate selected from the group  
4 consisting of PS80JJ1, having the identifying characteristics of NRRL B-18679; PS149B1,  
5 having the identifying characteristics of NRRL B-21553; and PS167H2, having the identifying  
6 characteristics of NRRL B-21554.

1           6. The isolated polynucleotide, according to claim 5, wherein said nucleotide sequence  
2 encodes a toxin of approximately 40-50 kDa.

1           7. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2 active against a non-mammalian pest wherein a portion of said nucleotide sequence can be  
3 amplified by PCR using a primer pair selected from the following group:

4           SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp; SEQ ID NOS. 20 and  
5 25 to produce a fragment of about 594 bp; SEQ ID NOS. 21 and 24 to produce a fragment of  
6 about 471 bp; and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp.

1           8. The isolated polynucleotide, according to claim 7, wherein said nucleotide sequence  
2 encodes a toxin of approximately 40-50 kDa.

1           9. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2 active against a non-mammalian pest wherein said toxin comprises a pesticidal portion of an  
3 amino acid sequence shown in the group selected from SEQ ID NO. 30, SEQ ID NO. 34, and  
4 SEQ ID NO. 39.

1           10. The isolated polynucleotide, according to claim 9, wherein said nucleotide sequence  
2 encodes a toxin which comprises a pesticidal portion of the consensus sequence shown in Figure  
3 1.

1           11. The isolated polynucleotide, according to claim 9, wherein said nucleotide sequence  
2 encodes a toxin of approximately 40-50 kDa.

1           12. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2 active against a non-mammalian pest wherein said toxin comprises an amino acid sequence  
3 which has at least about 75% homology with a pesticidal portion of an amino acid sequence  
4 selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ  
5 ID NO. 38, and SEQ ID NO. 43.

1           13. The isolated polynucleotide, according to claim 12, wherein said nucleotide  
2 sequence encodes a toxin which comprises an amino acid sequence which has at least about 80%  
3 homology with a pesticidal portion of an amino acid sequence selected from the group consisting  
4 of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43.

1           14. The isolated polynucleotide, according to claim 12, wherein said nucleotide  
2           sequence encodes a toxin which comprises an amino acid sequence which has at least about 90%  
3           homology with a pesticidal portion of an amino acid sequence selected from the group consisting  
4           of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43.

1           15. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           which is active against a non-mammalian pest, wherein said nucleotide sequence is from a  
3           *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1, having the  
4           identifying characteristics of NRRL B-18679; PS149B1, having the identifying characteristics  
5           of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554; and  
6           mutants thereof which retain pesticidal activity.

1           16. The isolated polynucleotide, according to claim 15, wherein said toxin is  
2           approximately 40-50 kDa.

1           17. The isolated polynucleotide, according to claim 15, wherein said toxin is  
2           approximately 10-15 kDa.

1           18. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           active against a non-mammalian pest wherein said nucleotide sequence hybridizes under  
3           stringent conditions with a nucleotide sequence selected from the group consisting of: DNA  
4           which encodes SEQ ID NO. 3; DNA which encodes SEQ ID NO. 5; and DNA which encodes  
5           SEQ ID NO. 7.

1           19. The isolated polynucleotide, according to claim 18, wherein said nucleotide  
2           sequence encodes a toxin of about 10-15 kDa.

1           20. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           active against a non-mammalian pest wherein said toxin immunoreacts with an antibody to an  
3           approximately 10-15 kDa toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate  
4           selected from the group consisting of PS80JJ1, having the identifying characteristics of NRRL  
5           B-18679; PS149B1, having the identifying characteristics of NRRL B-21553; and PS167H2,  
6           having the identifying characteristics of NRRL B-21554.

1           21. The isolated polynucleotide, according to claim 20, wherein said nucleotide  
2           sequence encodes a toxin of approximately 10-15 kDa.

1           22. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           active against a non-mammalian pest wherein a portion of said nucleotide sequence can be  
3           amplified by PCR using the primer pair of SEQ ID NO. 29 and SEQ ID NO. 33.

1           23. The isolated polynucleotide, according to claim 22, wherein said nucleotide  
2           sequence encodes a toxin of approximately 10-15 kDa.

1           24. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           active against a non-mammalian pest wherein said toxin comprises a pesticidal portion of an  
3           amino acid sequence selected from the group consisting of SEQ ID NO. 32, SEQ ID NO. 36, and  
4           SEQ ID NO. 41.

1           25. The isolated polynucleotide, according to claim 24, wherein said toxin comprises  
2           the amino acid sequence shown in SEQ ID NO. 32.

1           26. The isolated polynucleotide, according to claim 24, wherein said nucleotide  
2           sequence encodes a toxin of approximately 10-15 kDa.

1           27. An isolated polynucleotide comprising a nucleotide sequence which encodes a toxin  
2           active against a non-mammalian pest wherein said toxin comprises an amino acid sequence  
3           which has at least about 75% homology with an amino acid sequence selected from the group  
4           consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO.  
5           32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

1           28. The isolated polynucleotide, according to claim 27, wherein said nucleotide  
2           sequence encodes a toxin which comprises an amino acid sequence which has at least about 80%  
3           homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 3,  
4           SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ  
5           ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

1           29. The isolated polynucleotide, according to claim 27, wherein said nucleotide  
2 sequence encodes a toxin which comprises an amino acid sequence which has at least about 90%  
3 homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 3,  
4 SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ  
5 ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

1           30. A purified toxin active against a non-mammalian pest, wherein said toxin is  
2 encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide  
3 sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 2; DNA  
4 which encodes SEQ ID NO. 4; DNA which encodes SEQ ID NO. 6; SEQ ID NO. 8; SEQ ID  
5 NO. 10; DNA which encodes SEQ ID NO. 11; SEQ ID NO. 12; DNA which encodes SEQ ID  
6 NO. 13; SEQ ID NO. 14; DNA which encodes SEQ ID NO. 15; DNA which encodes SEQ ID  
7 NO. 16; DNA which encodes SEQ ID NO. 17; DNA which encodes SEQ ID NO. 18; DNA  
8 which encodes SEQ ID NO. 19; SEQ ID NO. 20; SEQ ID NO. 21; SEQ ID NO. 22; SEQ ID NO.  
9 23; SEQ ID NO. 24; SEQ ID NO. 25; SEQ ID NO. 26; SEQ ID NO. 27; DNA which encodes  
10 a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38,  
11 SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43.

1           31. The purified toxin, according to claim 30, wherein said toxin does not have the  
2 amino acid sequence shown in SEQ ID NO. 11.

1           32. The purified toxin, according to claim 31, wherein said toxin is encoded by a  
2 nucleotide sequence which hybridizes with DNA which encodes SEQ ID NO. 2.

1           33. The purified toxin, according to claim 31, which is encoded by DNA which  
2 hybridizes with SEQ ID NO. 10.

1           34. The purified toxin, according to claim 31, having a molecular weight of  
2 approximately 40-50 kDa.

1           35. A purified toxin active against a non-mammalian pest, wherein said toxin  
2 immunoreacts with an antibody to an approximately 40-50 kDa toxin, or a fragment thereof,  
3 from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1, having the

4 identifying characteristics of NRRL B-18679; PS149B1, having the identifying characteristics  
5 of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554.

1 36. The purified toxin, according to claim 35, wherein said toxin does not have the  
2 amino acid sequence shown in SEQ ID NO. 11.

1 37. The purified toxin, according to claim 36, having a molecular weight of about 40-50  
2 kDa.

1 38. A purified toxin having activity against a non-mammalian pest, wherein said toxin  
2 is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be  
3 amplified by PCR using a primer pair selected from the following group:

4 SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp; SEQ ID NOS. 20 and  
5 25 to produce a fragment of about 594 bp; SEQ ID NOS. 21 and 24 to produce a fragment of  
6 about 471 bp; and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp.

1 39. The purified toxin, according to claim 38, wherein said toxin does not have the  
2 amino acid sequence shown in SEQ ID NO. 11.

1 40. The purified toxin, according to claim 39, having a molecular weight of about 40-50  
2 kDa.

1 41. A purified toxin active against a non-mammalian pest, wherein said toxin comprises  
2 a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28.

1 42. The purified toxin, according to claim 41, wherein said toxin does not have the  
2 amino acid sequence shown in SEQ ID NO. 11.

1 43. The purified toxin, according to claim 42, wherein said toxin comprises a pesticidal  
2 portion of the consensus sequence in Figure 1.

1 44. The purified toxin, according to claim 42, having a molecular weight of  
2 approximately 40-50 kDa.

1           45. A purified toxin active against a non-mammalian pest, wherein said toxin comprises  
2 an amino acid sequence which has at least about 75% homology with a pesticidal portion of an  
3 amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13,  
4 SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43.

1           46. The purified toxin, according to claim 45, wherein said toxin does not have the  
2 amino acid sequence shown in SEQ ID NO. 11.

1           47. The purified toxin, according to claim 46, which comprises an amino acid sequence  
2 which has at least about 80% homology with a pesticidal portion of an amino acid sequence  
3 selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, and SEQ ID NO. 15,  
4 SEQ ID NO. 38, and SEQ ID NO. 43.

1           48. The purified toxin, according to claim 46, which comprises an amino acid sequence  
2 which has at least about 90% homology with a pesticidal portion of an amino acid sequence  
3 selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ  
4 ID NO. 38, and SEQ ID NO. 43.

1           49. The purified toxin, according to claim 46, having a molecular weight of  
2 approximately 40-50 kDa.

1           50. A purified toxin active against a non-mammalian pest, wherein said toxin is  
2 encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide  
3 sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 3; DNA  
4 which encodes SEQ ID NO. 5; and DNA which encodes SEQ ID NO. 7.

1           51. The purified toxin, according to claim 50, having a molecular weight of  
2 approximately 10-15 kDa.

1           52. A purified toxin active against a non-mammalian pest, wherein said toxin  
2 immunoreacts with an antibody to an approximately 10-15 kDa toxin, or a fragment thereof,  
3 from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1, having the  
4 identifying characteristics of NRRL B-18679; PS149B1, having the identifying characteristics  
5 of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554.

1           53. The purified toxin, according to claim 52, having a molecular weight of about 10-15  
2     kDa.

1           54. A purified toxin having activity against a non-mammalian pest, wherein said toxin  
2     is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be  
3     amplified by PCR using the primer pair of SEQ ID NO. 29 and SEQ ID NO. 33.

1           55. The purified toxin, according to claim 54, having a molecular weight of about 10-15  
2     kDa.

1           56. A purified toxin active against a non-mammalian pest, wherein said toxin comprises  
2     a pesticidal portion of an amino acid sequence selected from the group consisting of SEQ ID  
3     NO. 32, SEQ ID NO. 36, and SEQ ID NO. 41.

1           57. The purified toxin, according to claim 56, wherein said toxin comprises the amino  
2     acid sequence shown in SEQ ID NO. 32.

1           58. The purified toxin, according to claim 56, having a molecular weight of  
2     approximately 10-15 kDa.

1           59. A purified toxin active against a non-mammalian pest, wherein said toxin comprises  
2     an amino acid sequence which has at least about 75% homology with an amino acid sequence  
3     selected from the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal  
4     portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of  
5     SEQ ID NO. 41.

1           60. The purified toxin, according to claim 59, which comprises an amino acid sequence  
2     which has at least about 80% homology with an amino acid sequence selected from the group  
3     consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO.  
4     32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

1           61. The purified toxin, according to claim 59, which comprises an amino acid sequence  
2     which has at least about 90% homology with an amino acid sequence selected from the group

consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

62. A biologically pure culture of a *Bacillus thuringiensis* isolate selected from the group consisting of PS149B1, having the identifying characteristics of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554; and mutants thereof which retain pesticidal activity.

63. The biologically pure culture, according to claim 62, wherein said *Bacillus thuringiensis* isolate is PS149B1, having the identifying characteristics of NRRL B-21553.

64. The biologically pure culture, according to claim 62, wherein said *Bacillus thuringiensis* isolate is PS167H2, having the identifying characteristics of NRRL B-21554.

65. A composition of matter for controlling coleopterans comprising a *Bacillus thuringiensis* isolate selected from the group consisting of PS149B1, having the identifying characteristics of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554; and mutants thereof which retain activity against coleopterans, in association with an agricultural carrier appropriate for use in controlling coleopterans.

66. A method for controlling a non-mammalian pest comprising contacting said pest with a pesticidal amount of a *Bacillus thuringiensis* isolate, or a toxin of said *Bacillus thuringiensis* isolate, wherein said isolate is selected from the group consisting of PS149B1, having the identifying characteristics of NRRL B-21553; and PS167H2, having the identifying characteristics of NRRL B-21554; and mutants thereof which retain pesticidal activity.

67. The method, according to claim 66, wherein said *Bacillus thuringiensis* isolate is PS149B1, having the identifying characteristics of NRRL B-21553.

68. The method, according to claim 66, wherein said *Bacillus thuringiensis* isolate is PS167H2, having the identifying characteristics of NRRL B-21554.

69. A method for controlling a non-mammalian pest which comprises contacting said pest with a pesticidal amount of a *Bacillus thuringiensis* toxin wherein said toxin has a characteristic selected from the group consisting of:

- (a) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO. 4, DNA which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which encodes SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO. 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25, SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38, SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (b) said toxin immunoreacts with an antibody to an approximately 40-50 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (c) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using a primer pair selected from the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (d) said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;

- 34 (e) said toxin comprises an amino acid sequence which has at least about 75%  
35 homology with an amino acid sequence selected from the group consisting of  
36 SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ  
37 ID NO. 43; and wherein said toxin does not have the amino acid sequence  
38 shown in SEQ ID NO. 11;
- 39 (f) said toxin is encoded by a nucleotide sequence which hybridizes under stringent  
40 conditions with a nucleotide sequence selected from the group consisting of  
41 DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO. 5, and  
42 DNA which encodes SEQ ID NO. 7;
- 43 (g) said toxin immunoreacts with an antibody to an approximately 10-15 kDa  
44 pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate  
45 selected from the group consisting of PS80JJ1 having the identifying  
46 characteristics of NRRL B-18679, PS149B1 having the identifying  
47 characteristics of NRRL B-21553, and PS167H2 having the identifying  
48 characteristics of NRRL B-21554;
- 49 (h) said toxin is encoded by a nucleotide sequence wherein a portion of said  
50 nucleotide sequence can be amplified by PCR using the primer pair of SEQ ID  
51 NO. 29 and SEQ ID NO. 33;
- 52 (i) said toxin comprises a pesticidal portion of an amino acid sequence selected  
53 from the group consisting of SEQ ID NO. 32, SEQ ID NO. 36, and SEQ ID NO.  
54 41; and
- 55 (j) said toxin comprises an amino acid sequence which has at least about 75%  
56 homology with an amino acid sequence selected from the group consisting of  
57 SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID  
58 NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of  
59 sequence IDS NO. 41.

1 70. The method, according to claim 69, wherein the full length of said toxin is  
2 approximately 40-50 kDa.

1 71. The method, according to claim 69, wherein said toxin is encoded by a nucleotide  
2 sequence which hybridizes under stringent conditions with a nucleotide sequence selected from  
3 the group consisting of: DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO.  
4 4, DNA which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which encodes

SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO. 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25, SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38, SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

72. The method, according to claim 69, wherein said toxin immunoreacts with an antibody to an approximately 40-50 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

73. The method, according to claim 69, wherein said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using a primer pair selected from the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

74. The method, according to claim 69, wherein said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

75. The method, according to claim 69, wherein said toxin comprises an amino acid sequence which has at least about 75% homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

1           76. The method, according to claim 69, wherein the full length of said toxin is  
2 approximately 10-15 kDa.

1           77. The method, according to claim 69, wherein said toxin is encoded by a nucleotide  
2 sequence which hybridizes under stringent conditions with a nucleotide sequence selected from  
3 the group consisting of DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO.  
4 5, and DNA which encodes SEQ ID NO. 7.

1           78. The method, according to claim 69, wherein said toxin immunoreacts with an  
2 antibody to an approximately 10-15 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus*  
3 *thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying  
4 characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-  
5 21553, and PS167H2 having the identifying characteristics of NRRL B-21554

1           79. The method, according to claim 69, wherein said toxin is encoded by a nucleotide  
2 sequence wherein a portion of said nucleotide sequence can be amplified by PCR using the  
3 primer pair of SEQ ID NO. 29 and SEQ ID NO. 33.

1           80. The method, according to claim 69, wherein said toxin comprises a pesticidal  
2 portion of an amino acid sequence selected from the group consisting of SEQ ID NO. 32, SEQ  
3 ID NO. 36, and SEQ ID NO. 41.

1           81. The method, according to claim 69, wherein said toxin comprises an amino acid  
2 sequence which has at least about 75% homology with an amino acid sequence selected from  
3 the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of  
4 SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS  
5 NO. 41.

1           82. The method, according to claim 69, wherein said pest is an insect.

1           83. The method, according to claim 82, wherein said insect is a coleopteran.

1           84. The method, according to claim 82, wherein said insect is a lepidopteran.

1 85. The method, according to claim 69, wherein said pest is a mite.

1 86. The method, according to claim 69, wherein said pest is corn rootworm.

1 87. The method, according to claim 69, wherein said toxin is encoded by DNA which  
2 hybridizes with SEQ ID NO. 2, SEQ ID NO. 10, SEQ ID NO. 37, or SEQ ID NO. 42, and  
3 wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11.

1 88. The method, according to claim 69, wherein said toxin comprises the consensus  
2 sequence shown in Figure 1 and wherein said toxin does not have the amino acid sequence  
3 shown in SEQ ID NO. 11.

1 89. The method, according to claim 69, wherein said toxin comprises an amino acid  
2 sequence which has at least about 75% homology with a pesticidal portion of an amino acid  
3 sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO.  
4 15, SEQ ID NO. 38, and SEQ ID NO. 43.

1 90. The method, according to claim 69, wherein said toxin comprises an amino acid  
2 sequence which has at least about 90% homology with a pesticidal portion of an amino acid  
3 sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO.  
4 15, SEQ ID NO. 38, and SEQ ID NO. 43.

1 91. The method, according to claim 69, wherein said toxin comprises the consensus  
2 sequence shown in Figure 1.

1 92. The method, according to claim 69, wherein said toxin comprises an amino acid  
2 sequence which has at least about 75% homology with an amino acid sequence selected from  
3 the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of  
4 SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS  
5 NO. 41.

1 93. The method, according to claim 69, wherein said toxin comprises an amino acid  
2 sequence which has at least about 90% homology with an amino acid sequence selected from

the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

94. A method for controlling a non-mammalian pest wherein said method comprises contacting said pest with a first toxin wherein said toxin has a characteristic selected from the group consisting of:

- (a) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO. 4, DNA which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which encodes SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO. 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25, SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38, SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (b) said toxin immunoreacts with an antibody to an approximately 40-50 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (c) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using a primer pair selected from the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;

- (d) said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (e) said toxin comprises an amino acid sequence which has at least about 75% homology with a pesticidal portion of an amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- and further comprising contacting said pest with a second toxin having a characteristic selected from the group consisting of:
- (f) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO. 5, and DNA which encodes SEQ ID NO. 7;
- (g) said toxin immunoreacts with an antibody to an approximately 10-15 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554;
- (h) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using the primer pair of SEQ ID NO. 29 and SEQ ID NO. 33;
- (i) said toxin comprises a pesticidal portion of an amino acid sequence selected from the group SEQ ID NO. 32, SEQ ID NO. 36, and SEQ ID NO. 41; and
- (j) said toxin comprises an amino acid sequence which has at least about 75% homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

95. The process, according to claim 94, wherein said pest is selected from the group consisting of insects and mites.

1 96. The process, according to claim 95, wherein said pest is a coleopteran.

1 97. The process, according to claim 94, wherein said first toxin has a full length of  
2 about 40-50 kDa and said second toxin has a full length of about 10-15 kDa.

1 98. A recombinant host transformed to express a toxin having activity against a non-  
2 mammalian pest wherein said toxin has at least one characteristic selected from the group  
3 consisting of:

- 4 (a) said toxin is encoded by a nucleotide sequence which hybridizes under stringent  
5 conditions with a nucleotide sequence selected from the group consisting of:  
6 DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO. 4, DNA  
7 which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which  
8 encodes SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO.  
9 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes  
10 SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes  
11 SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ  
12 ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25,  
13 SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of  
14 SEQ ID NO. 28, SEQ ID NO. 37, DNA which encodes SEQ ID NO. 38, SEQ  
15 ID NO. 42, and DNA which encodes SEQ ID NO. 43; and wherein said toxin  
16 does not have the amino acid sequence shown in SEQ ID NO. 11;
- 17 (b) said toxin immunoreacts with an antibody to an approximately 40-50 kDa  
18 pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate  
19 selected from the group consisting of PS80JJ1 having the identifying  
20 characteristics of NRRL B-18679, PS149B1 having the identifying  
21 characteristics of NRRL B-21553, and PS167H2 having the identifying  
22 characteristics of NRRL B-21554, and wherein said toxin does not have the  
23 amino acid sequence shown in SEQ ID NO. 11;
- 24 (c) said toxin is encoded by a nucleotide sequence wherein a portion of said  
25 nucleotide sequence can be amplified by PCR using a primer pair selected from  
26 the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about  
27 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ  
28 ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS.

- 21 and 25 to produce a fragment of about 580 bp, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (d) said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (e) said toxin comprises an amino acid sequence which has at least about 75% homology with a pesticidal portion of an amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (f) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO. 5, and DNA which encodes SEQ ID NO. 7;
- (g) said toxin immunoreacts with an antibody to an approximately 10-15 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554;
- (h) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using the primer pair of SEQ ID NO. 29 and SEQ ID NO. 33;
- (i) said toxin comprises a pesticidal portion of an amino acid sequence selected from the group consisting of SEQ ID NO. 32, SEQ ID NO. 36, and SEQ ID NO. 41; and
- (j) said toxin comprises an amino acid sequence which has at least about 75% homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of sequence IDS NO. 41.

99. The recombinant host, according to claim 98, wherein said host expresses a first toxin which has a characteristic selected from the group consisting of:

- (a) said toxin is encoded by a nucleotide sequence which hybridizes under stringent conditions with a nucleotide sequence selected from the group consisting of: DNA which encodes SEQ ID NO. 2, DNA which encodes SEQ ID NO. 4, DNA which encodes SEQ ID NO. 6, SEQ ID NO. 8, SEQ ID NO. 10, DNA which encodes SEQ ID NO. 11, SEQ ID NO. 12, DNA which encodes SEQ ID NO. 13, SEQ ID NO. 14, DNA which encodes SEQ ID NO. 15, DNA which encodes SEQ ID NO. 16, DNA which encodes SEQ ID NO. 17, DNA which encodes SEQ ID NO. 18, DNA which encodes SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, SEQ ID NO. 22, SEQ ID NO. 23, SEQ ID NO. 24, SEQ ID NO. 25, SEQ ID NO. 26, SEQ ID NO. 27, DNA which encodes a pesticidal portion of SEQ ID NO. 28, SEQ ID NO. 39, DNA which encodes SEQ ID NO. 38, SEQ ID NO. 42, and DNA which encodes SEQ ID NO. 43; and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (b) said toxin immunoreacts with an antibody to an approximately 40-50 kDa pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate selected from the group consisting of PS80JJ1 having the identifying characteristics of NRRL B-18679, PS149B1 having the identifying characteristics of NRRL B-21553, and PS167H2 having the identifying characteristics of NRRL B-21554, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (c) said toxin is encoded by a nucleotide sequence wherein a portion of said nucleotide sequence can be amplified by PCR using a primer pair selected from the group consisting of SEQ ID NOS. 20 and 24 to produce a fragment of about 495 bp, SEQ ID NOS. 20 and 25 to produce a fragment of about 594 bp, SEQ ID NOS. 21 and 24 to produce a fragment of about 471 bp, and SEQ ID NOS. 21 and 25 to produce a fragment of about 580 bp, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11;
- (d) said toxin comprises a pesticidal portion of the amino acid sequence shown in SEQ ID NO. 28, and wherein said toxin does not have the amino acid sequence shown in SEQ ID NO. 11; and
- (e) said toxin comprises an amino acid sequence which has at least about 75% homology with an amino acid sequence selected from the group consisting of SEQ ID NO. 11, SEQ ID NO. 13, SEQ ID NO. 15, SEQ ID NO. 38, and SEQ

36 ID NO. 43; and wherein said toxin does not have the amino acid sequence  
37 shown in SEQ ID NO. 11;

38 and said host expresses a second toxin having a characteristic selected from the group consisting  
39 of:

- 40 (f) said toxin is encoded by a nucleotide sequence which hybridizes under stringent  
41 conditions with a nucleotide sequence selected from the group consisting of  
42 DNA which encodes SEQ ID NO. 3, DNA which encodes SEQ ID NO. 5, and  
43 DNA which encodes SEQ ID NO. 7;
- 44 (g) said toxin immunoreacts with an antibody to an approximately 10-15 kDa  
45 pesticidal toxin, or a fragment thereof, from a *Bacillus thuringiensis* isolate  
46 selected from the group consisting of PS80JJ1 having the identifying  
47 characteristics of NRRL B-18679, PS149B1 having the identifying  
48 characteristics of NRRL B-21553, and PS167H2 having the identifying  
49 characteristics of NRRL B-21554;
- 50 (h) said toxin is encoded by a nucleotide sequence wherein a portion of said  
51 nucleotide sequence can be amplified by PCR using the primer pair of SEQ ID  
52 NO. 29 and SEQ ID NO. 33;
- 53 (i) said toxin comprises a pesticidal portion of an amino acid sequence selected  
54 from the group consisting of SEQ ID NO. 32, SEQ ID NO. 36, and SEQ ID NO.  
55 41; and
- 56 (j) said toxin comprises an amino acid sequence which has at least about 75%  
57 homology with an amino acid sequence selected from the group consisting of  
58 SEQ ID NO. 3, SEQ ID NO. 5, SEQ ID NO. 7, pesticidal portions of SEQ ID  
59 NO. 32, pesticidal portions of SEQ ID NO. 36, and pesticidal portions of  
60 sequence IDS NO. 41.

1 100. The recombinant host, according to claim 98, wherein said host is selected from  
2 the group consisting of plants, yeasts, and bacteria.

1 101. The isolated polynucleotide, according to claim 1, wherein said nucleotide  
2 sequence hybridizes with SEQ ID NO. 37.

1 102. The isolated polynucleotide, according to claim 1, wherein said nucleotide  
2 sequence hybridizes with SEQ ID NO. 42.

1           103. The purified toxin, according to claim 31, which is encoded by DNA which  
2 hybridizes with SEQ ID NO. 37.

1           104. The purified toxin, according to claim 31, which is encoded by DNA which  
2 hybridizes with SEQ ID NO. 42.

1           105. The isolated polynucleotide, according to claim 24, wherein said toxin comprises  
2 the amino acid sequence shown in SEQ ID NO. 36.

1           106. The isolated polynucleotide, according to claim 24, wherein said toxin comprises  
2 the amino acid sequence shown in SEQ ID NO. 41.

1           107. The purified toxin, according to claim 56, wherein said toxin comprises the amino  
2 acid sequence shown in SEQ ID NO. 36.

1           108. The purified toxin, according to claim 56, wherein said toxin comprises the amino  
2 acid sequence shown in SEQ ID NO. 41.

1           109. An isolated polynucleotide comprising a nucleotide sequence which encodes an  
2 approximately 10-15 kDa 80JJ1 toxin active against non-mammalian pests, wherein said  
3 nucleotide sequence has been optimized for expression in plants.

1           110. The isolated polynucleotide, according to claim 109, wherein said polynucleotide  
2 comprises the sequence shown in SEQ ID NO. 44.

1           111. An isolated polynucleotide comprising a nucleotide sequence which encodes an  
2 approximately 40-50 kDa 80JJ1 toxin active against non-mammalian pests, wherein said  
3 nucleotide sequence has been optimized for expression in plants.

1           112. The isolated polynucleotide, according to claim 111, wherein said polynucleotide  
2 comprises the sequence shown in SEQ ID NO. 45.

1/2

## FIG. 1

	1		50
{149b145k}	.....GLYAA	TYLSLDDSGV	SLMNKNDDDI DDYNLKWFLF
{167h245k}	.....HAA	TYLSLDDSGV	SLMNKNDDDI DDYNLKWFLF
{80jj145k}	MLDTNKVYEI	TYLSLDDSGV	SLMSKKDEDI DDYNLKWFLF
Consensus	-----	TYLSLDDSGV	SLM-K-D-DI DDYNL-WELF
	51		100
{149b145k}	PIDDDQYIIT	SYAANNCKVW	NVNNDKINVS TYSSSNSIQK WQIKANGSSY
{167h245k}	PIDDNQYIIT	SYAANNCKVW	NVNNDKINVS TYSSSNSIQK WQIKANASSY
{80jj145k}	PIDNNQYIIT	SYGANNCKVW	NVKNDKINVS TYSSSNSVQK WQIKAKDSSY
Consensus	PID--QYIIT	SY-ANNCKVW	NV-NDKINVS TYSSSNS-QK WQIKA--SSY
	101		150
{149b145k}	VIQSDNGKVL	TAGTGQALGL	IRLTDESSNN PNQQWNLTSV QTIQLPQKPI
{167h245k}	VIQSNNGKVL	TAGTGQSLGL	IRLTDESPDN PNQQWNLTPV QTIQLPPKPT
{80jj145k}	IIQSDNGKVL	TAGVGQSLGI	VRLTDEFPEN SNQQWNLTPV QTIQLPQKPK
Consensus	-IQS-NGKVL	TAG-GQ-LG-	-RLTDE---N -NQQWNL-T-V QTIQLP-KP-
	151		200
{149b145k}	IDTKLKDYPK	YSPGTGNIDNG	TSPQLMGWTL VPCIMVNDPN IDKNTQIKTT
{167h245k}	IDTKLKDYPK	YSQTGNIDKG	TPPQLMGWTL IPCIMVNDPN IDKNTQIKTT
{80jj145k}	IDEKLKDHP	YSETGNINPK	TTPQLMGWTL VPCIMVNDPK IDKNTQIKTT
Consensus	ID-KLKD-P-	YS-TGNI---	T-PQLMGWTL -PCIMVND-- IDKNTQIKTT
	201		250
{149b145k}	PYYILKKYQY	WQRAVGSNVA	LRPHEKKSXT YEWGTEIDQK TTIINTLGFEQ
{167h245k}	PYYILKKYQY	WQRAVGSNVA	LRPHEKKSXT YEWGTEIDQK TTIINTLGFEQ
{80jj145k}	PYYIFKKYKY	WNLAKGSNVS	LLPHQKRSYD YEWGTEKQK TTIINTVGLQ
Consensus	PYYI-KKY-Y	W--A-GSNV-	L-PH-K-SY- YEWGTE--QK TTIINT-G-Q
	251		300
{149b145k}	INIDSGMKFD	IPEVGGGTDE	IKTQLNEELK IEYSHETKIM EKY.....
{167h245k}	INIDSGMKFD	IPEVGGGTDE	IKTQLNEELK IEYSRETAKIM EKY.....
{80jj145k}	INIDSGMKFE	VPEVGGGTED	IKTQLTEELK VEYSRETAKIM TKYQEHSEID
Consensus	INIDSGMKF-	-PEVGGGT--	IKTQL-EEELK -EYS-ETKIM -KY-----
	301		350
{149b145k}	.....	.....	.....
{167h245k}	.....	.....	.....
{80jj145k}	NPTNQPMNSI	GLLIYTSLEL	YRYNGTEIKI MDISTSDHDT YTLTSYPNHK
Consensus	-----	-----	-----
	351		386
{149b145k}	.....	.....	.....
{167h245k}	.....	.....	.....
{80jj145k}	EALLLTNHS	YEEVEETIKI	PKHTLIKLRK HYFKK.
Consensus	-----	-----	-----

2/2

FIG. 2

